

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷: C12N 9/10, 9/02, 15/82, 15/29, 15/63, 15/83, C12Q 1/48, 1/26, 1/68, G01N 33/50	A3	(11) International Publication Number: WO 00/32757 (43) International Publication Date: 8 June 2000 (08.06.00)
(21) International Application Number: PCT/US99/28588 (22) International Filing Date: 2 December 1999 (02.12.99) (30) Priority Data: 60/110,781 3 December 1998 (03.12.98) US (71) Applicant (for all designated States except US): E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): CAHOON, Rebecca, E. [US/US]; 2331 West 18th Street, Wilmington, DE 19806 (US). COUGHLAN, Sean, J. [US/US]; 108 Stratton Drive, Hockessin, DE 19707 (US). MIAO, Guo-Hua [US/US]; 202 Cherry Blossom Place, Hockessin, DE 19707 (US). RAFALSKI, J., Antoni [US/US]; 2028 Longcome Drive, Wilmington, DE 19810 (US). (74) Agent: FEULNER, Gregory, J.; E.I. du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		(81) Designated States: AE, AL, AU, BA, BB, BG, BR, CA, CN, CR, CU, CZ, DM, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 23 November 2000 (23.11.00)
(54) Title: PLANT VITAMIN E BIOSYNTHETIC ENZYMES (57) Abstract <p>This invention relates to an isolated nucleic acid fragment encoding a vitamin E biosynthetic enzyme. The invention also relates to the construction of a chimeric gene encoding all or a portion of the vitamin E biosynthetic enzyme, in sense or antisense orientation, wherein expression of the chimeric gene results in production of altered levels of the vitamin E biosynthetic enzyme in a transformed host cell.</p> <pre> SEQ ID NO:39 NVTNVPKHAL-----FLAFICYTS SEQ ID NO:40 NKATLAAPSLTSL-P-----YTHNS-FGSKSLFSPSSSSVS-NTTTRGV SEQ ID NO:38 NNN-----SKNPA-PLTFLHLDAAFPFPPFIDZKAARVPPVPLPLPAR-XLAAPOCV SEQ ID NO:28 NATVRI-----PTISCIIRHTFSQSPRTFARIRVGFNPAIAXA-ASSGEI SEQ ID NO:08 NATVRI-----PTISCIIRHTFSQSPRTFARIRVGFNPAIAXA-ASSGEI SEQ ID NO:22 NAAALLNCSOSS-SLAACRGEVIRAPSH-VFNNRRLARAVYSL--R-PHASTA- SEQ ID NO:24 AR--NPTGLALHPLLCTSSMLCASAPACLLKNNRRKSSRHTLYAVKAAHP SEQ ID NO:26 FNN-----GH----- 1 60 120 180 240 300 360 380 390 400 410 420 430 440 450 460 470 480 490 500 510 520 530 540 550 560 570 580 590 600 610 620 630 640 650 660 670 680 690 700 710 720 730 740 750 760 770 780 790 800 810 820 830 840 850 860 870 880 890 900 910 920 930 940 950 960 970 980 990 1000 1010 1020 1030 1040 1050 1060 1070 1080 1090 1100 1110 1120 1130 1140 1150 1160 1170 1180 1190 1200 1210 1220 1230 1240 1250 1260 1270 1280 1290 1300 1310 1320 1330 1340 1350 1360 1370 1380 1390 1400 1410 1420 1430 1440 1450 1460 1470 1480 1490 1500 1510 1520 1530 1540 1550 1560 1570 1580 1590 1600 1610 1620 1630 1640 1650 1660 1670 1680 1690 1700 1710 1720 1730 1740 1750 1760 1770 1780 1790 1800 1810 1820 1830 1840 1850 1860 1870 1880 1890 1900 1910 1920 1930 1940 1950 1960 1970 1980 1990 2000 2010 2020 2030 2040 2050 2060 2070 2080 2090 2100 2110 2120 2130 2140 2150 2160 2170 2180 2190 2200 2210 2220 2230 2240 2250 2260 2270 2280 2290 2300 2310 2320 2330 2340 2350 2360 2370 2380 2390 2400 2410 2420 2430 2440 2450 2460 2470 2480 2490 2500 2510 2520 2530 2540 2550 2560 2570 2580 2590 2600 2610 2620 2630 2640 2650 2660 2670 2680 2690 2700 2710 2720 2730 2740 2750 2760 2770 2780 2790 2800 2810 2820 2830 2840 2850 2860 2870 2880 2890 2900 2910 2920 2930 2940 2950 2960 2970 2980 2990 3000 3010 3020 3030 3040 3050 3060 3070 3080 3090 3100 3110 3120 3130 3140 3150 3160 3170 3180 3190 3200 3210 3220 3230 3240 3250 3260 3270 3280 3290 3300 3310 3320 3330 3340 3350 3360 3370 3380 3390 3400 3410 3420 3430 3440 3450 3460 3470 3480 3490 3500 3510 3520 3530 3540 3550 3560 3570 3580 3590 3600 3610 3620 3630 3640 3650 3660 3670 3680 3690 3700 3710 3720 3730 3740 3750 3760 3770 3780 3790 3800 3810 3820 3830 3840 3850 3860 3870 3880 3890 3900 3910 3920 3930 3940 3950 3960 3970 3980 3990 4000 4010 4020 4030 4040 4050 4060 4070 4080 4090 4100 4110 4120 4130 4140 4150 4160 4170 4180 4190 4200 4210 4220 4230 4240 4250 4260 4270 4280 4290 4300 4310 4320 4330 4340 4350 4360 4370 4380 4390 4400 4410 4420 4430 4440 4450 4460 4470 4480 4490 4500 4510 4520 4530 4540 4550 4560 4570 4580 4590 4600 4610 4620 4630 4640 4650 4660 4670 4680 4690 4700 4710 4720 4730 4740 4750 4760 4770 4780 4790 4800 4810 4820 4830 4840 4850 4860 4870 4880 4890 4900 4910 4920 4930 4940 4950 4960 4970 4980 4990 5000 5010 5020 5030 5040 5050 5060 5070 5080 5090 5100 5110 5120 5130 5140 5150 5160 5170 5180 5190 5200 5210 5220 5230 5240 5250 5260 5270 5280 5290 5300 5310 5320 5330 5340 5350 5360 5370 5380 5390 5400 5410 5420 5430 5440 5450 5460 5470 5480 5490 5500 5510 5520 5530 5540 5550 5560 5570 5580 5590 5600 5610 5620 5630 5640 5650 5660 5670 5680 5690 5700 5710 5720 5730 5740 5750 5760 5770 5780 5790 5800 5810 5820 5830 5840 5850 5860 5870 5880 5890 5900 5910 5920 5930 5940 5950 5960 5970 5980 5990 6000 6010 6020 6030 6040 6050 6060 6070 6080 6090 6100 6110 6120 6130 6140 6150 6160 6170 6180 6190 6200 6210 6220 6230 6240 6250 6260 6270 6280 6290 6300 6310 6320 6330 6340 6350 6360 6370 6380 6390 6400 6410 6420 6430 6440 6450 6460 6470 6480 6490 6500 6510 6520 6530 6540 6550 6560 6570 6580 6590 6600 6610 6620 6630 6640 6650 6660 6670 6680 6690 6700 6710 6720 6730 6740 6750 6760 6770 6780 6790 6800 6810 6820 6830 6840 6850 6860 6870 6880 6890 6900 6910 6920 6930 6940 6950 6960 6970 6980 6990 7000 7010 7020 7030 7040 7050 7060 7070 7080 7090 7100 7110 7120 7130 7140 7150 7160 7170 7180 7190 7200 7210 7220 7230 7240 7250 7260 7270 7280 7290 7300 7310 7320 7330 7340 7350 7360 7370 7380 7390 7400 7410 7420 7430 7440 7450 7460 7470 7480 7490 7500 7510 7520 7530 7540 7550 7560 7570 7580 7590 7600 7610 7620 7630 7640 7650 7660 7670 7680 7690 7700 7710 7720 7730 7740 7750 7760 7770 7780 7790 7800 7810 7820 7830 7840 7850 7860 7870 7880 7890 7900 7910 7920 7930 7940 7950 7960 7970 7980 7990 8000 8010 8020 8030 8040 8050 8060 8070 8080 8090 8100 8110 8120 8130 8140 8150 8160 8170 8180 8190 8200 8210 8220 8230 8240 8250 8260 8270 8280 8290 8300 8310 8320 8330 8340 8350 8360 8370 8380 8390 8400 8410 8420 8430 8440 8450 8460 8470 8480 8490 8500 8510 8520 8530 8540 8550 8560 8570 8580 8590 8600 8610 8620 8630 8640 8650 8660 8670 8680 8690 8700 8710 8720 8730 8740 8750 8760 8770 8780 8790 8800 8810 8820 8830 8840 8850 8860 8870 8880 8890 8900 8910 8920 8930 8940 8950 8960 8970 8980 8990 9000 9010 9020 9030 9040 9050 9060 9070 9080 9090 9100 9110 9120 9130 9140 9150 9160 9170 9180 9190 9200 9210 9220 9230 9240 9250 9260 9270 9280 9290 9300 9310 9320 9330 9340 9350 9360 9370 9380 9390 9400 9410 9420 9430 9440 9450 9460 9470 9480 9490 9500 9510 9520 9530 9540 9550 9560 9570 9580 9590 9600 9610 9620 9630 9640 9650 9660 9670 9680 9690 9700 9710 9720 9730 9740 9750 9760 9770 9780 9790 9800 9810 9820 9830 9840 9850 9860 9870 9880 9890 9900 9910 9920 9930 9940 9950 9960 9970 9980 9990 10000 10010 10020 10030 10040 10050 10060 10070 10080 10090 10100 10110 10120 10130 10140 10150 10160 10170 10180 10190 10200 10210 10220 10230 10240 10250 10260 10270 10280 10290 10300 10310 10320 10330 10340 10350 10360 10370 10380 10390 10400 10410 10420 10430 10440 10450 10460 10470 10480 10490 10500 10510 10520 10530 10540 10550 10560 10570 10580 10590 10600 10610 10620 10630 10640 10650 10660 10670 10680 10690 10700 10710 10720 10730 10740 10750 10760 10770 10780 10790 10800 10810 10820 10830 10840 10850 10860 10870 10880 10890 10900 10910 10920 10930 10940 10950 10960 10970 10980 10990 11000 11010 11020 11030 11040 11050 11060 11070 11080 11090 11100 11110 11120 11130 11140 11150 11160 11170 11180 11190 11200 11210 11220 11230 11240 11250 11260 11270 11280 11290 11300 11310 11320 11330 11340 11350 11360 11370 11380 11390 11400 11410 11420 11430 11440 11450 11460 11470 11480 11490 11500 11510 11520 11530 11540 11550 11560 11570 11580 11590 11600 11610 11620 11630 11640 11650 11660 11670 11680 11690 11700 11710 11720 11730 11740 11750 11760 11770 11780 11790 11800 11810 11820 11830 11840 11850 11860 11870 11880 11890 11900 11910 11920 11930 11940 11950 11960 11970 11980 11990 12000 12010 12020 12030 12040 12050 12060 12070 12080 12090 12100 12110 12120 12130 12140 12150 12160 12170 12180 12190 12200 12210 12220 12230 12240 12250 12260 12270 12280 12290 12300 12310 12320 12330 12340 12350 12360 12370 12380 12390 12400 12410 12420 12430 12440 12450 12460 12470 12480 12490 12500 12510 12520 12530 12540 12550 12560 12570 12580 12590 12600 12610 12620 12630 12640 12650 12660 12670 12680 12690 12700 12710 12720 12730 12740 12750 12760 12770 12780 12790 12800 12810 12820 12830 12840 12850 12860 12870 12880 12890 12900 12910 12920 12930 12940 12950 12960 12970 12980 12990 13000 13010 13020 13030 13040 13050 13060 13070 13080 13090 13100 13110 13120 13130 13140 13150 13160 13170 13180 13190 13200 13210 13220 13230 13240 13250 13260 13270 13280 13290 13300 13310 13320 13330 13340 13350 13360 13370 13380 13390 13400 13410 13420 13430 13440 13450 13460 13470 13480 13490 13500 13510 13520 13530 13540 13550 13560 13570 13580 13590 13600 13610 13620 13630 13640 13650 13660 13670 13680 13690 13700 13710 13720 13730 13740 13750 13760 13770 13780 13790 13800 13810 13820 13830 13840 13850 13860 13870 13880 13890 13900 13910 13920 13930 13940 13950 13960 13970 13980 13990 14000 14010 14020 14030 14040 14050 14060 14070 14080 14090 14100 14110 14120 14130 14140 14150 14160 14170 14180 14190 14200 14210 14220 14230 14240 14250 14260 14270 14280 14290 14300 14310 14320 14330 14340 14350 14360 14370 14380 14390 14400 14410 14420 14430 14440 14450 14460 14470 14480 14490 14500 14510 14520 14530 14540 14550 14560 14570 14580 14590 14600 14610 14620 14630 14640 14650 14660 14670 14680 14690 14700 14710 14720 14730 14740 14750 14760 14770 14780 14790 14800 14810 14820 14830 14840 14850 14860 14870 14880 14890 14900 14910 14920 14930 14940 14950 14960 14970 14980 14990 15000 15010 15020 15030 15040 15050 15060 15070 15080 15090 15100 15110 15120 15130 15140 15150 15160 15170 15180 15190 15200 15210 15220 15230 15240 15250 15260 15270 15280 15290 15300 15310 15320 15330 15340 15350 15360 15370 15380 15390 15400 15410 15420 15430 15440 15450 15460 15470 15480 15490 15500 15510 15520 15530 15540 15550 15560 15570 15580 15590 15600 15610 15620 15630 15640 15650 15660 15670 15680 15690 15700 15710 15720 15730 15740 15750 15760 15770 15780 15790 15800 15810 15820 15830 15840 15850 15860 15870 15880 15890 15900 15910 15920 15930 15940 15950 15960 15970 15980 15990 16000 16010 16020 16030 16040 16050 16060 16070 16080 16090 16100 16110 16120 16130 16140 16150 16160 16170 16180 16190 16200 16210 16220 16230 16240 16250 16260 16270 16280 16290 16300 16310 16320 16330 16340 16350 16360 16370 16380 16390 16400 16410 16420 16430 16440 16450 16460 16470 16480 16490 16500 16510 16520 16530 16540 16550 16560 16570 16580 16590 16600 16610 16620 16630 16640 16650 16660 16670 16680 16690 16700 16710 16720 16730 16740 16750 16760 16770 16780 16790 16800 16810 16820 16830 16840 16850 16860 16870 16880 16890 16900 16910 16920 16930 16940 16950 16960 16970 16980 16990 17000 17010 17020 17030 17040 17050 17060 17070 17080 17090 17100 17110 17120 17130 17140 17150 17160 17170 17180 17190 17200 17210 17220 17230 17240 17250 17260 17270 17280 17290 17300 17310 17320 17330 17340 17350 17360 17370 17380 17390 17400 17410 17420 17430 17440 17450 17460 17470 17480 17490 17500 17510 17520 17530 17540 17550 17560 17570 17580 17590 17600 17610 17620 17630 17640 17650 17660 17670 17680 17690 17700 17710 17720 17730 17740 17750 17760 17770 17780 17790 17800 17810 17820 17830 17840 17850 17860 17870 17880 17890 17900 17910 17920 17930 17940 17950 17960 17970 17980 17990 18000 18010 18020 18030 18040 18050 18060 18070 18080 18090 18100 18110 18120 18130 18140 18150 18160 18170 18180 18190 18200 18210 18220 18230 18240 18250 18260 18270 18280 18290 18300 18310 18320 18330 18340 18350 18360 18370 18380 18390 18400 18410 18420 18430 18440 18450 18460 18470 18480 18490 18500 18510 18520 18530 18540 18550 18560 18570 18580 18590 18600 18610 18620 18630 18640 18650 18660 18670 18680 18690 18700 18710 18720 18730 18740 18750 18760 18770 18780 18790 18800 18810 18820 18830 18840 18850 18860 18870 18880 18890 18900 18910 18920 18930 18940 18950 18960 18970 18980 18990 19000 19010 19020 19030 19040 19050 19060 19070 19080 19090 19100 19110 19120 19130 19140 19150 19160 19170 19180 19190 19200 19210 19220 19230 19240 19250 19260 19270 19280 19290 19300 19310 19320 19330 19340 19350 19360 19370 19380 19390 19400 19410 19420 19430 19440 19450 19460 19470 19480 19490 19500 19510 19520 19530 19540 19550 19560 19570 19580 19590 19600 19610 19620 19630 19640 19650 19660 19670 19680 19690 19700 19710 19720 19730 19740 19750 19760 19770 19780 19790 19800 19810 19820 19830 19840 19850 19860 19870 19880 19890 19900 19910 </pre>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

Internat : Application No
PCT/US 99/28588

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/10 C12N9/02 C12N15/82 C12N15/29 C12N15/63
C12N15/83 C12Q1/48 C12Q1/26 C12Q1/68 G01N33/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL [Online] Accession No. D64004, 4 October 1995 (1995-10-04) "Synechocystis sp. PCC6803 complete genome, delta(24)-sterol C-methyltransferase" XP002135563 see p. 22, l. 38 of present description: 1001725 corresponds to BAA10562 and D64004 the whole document</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	<p>1,3-11, 13-25</p>



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

12 April 2000

Date of mailing of the international search report

10 1 09 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Herrmann, K

INTERNATIONAL SEARCH REPORT

Internal Application No
PCT/US 99/28588

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BARTLEY, G.E. ET AL.: "Cloning of an Arabidopsis thaliana cDNA for p-hydroxyphenylpyruvate dioxygenase (Accession No. U89267)" PLANT PHYSIOL., vol. 113, 1997, pages 1465-1465, XP002135561 the whole document	
A	& DATABASE EMBL [Online] Accession No. U89267, 19 March 1997 (1997-03-19) "Arabidopsis thaliana p-hydroxyphenylpyruvate dioxygenase (HPD) mRNA, complete cds" the whole document	
A	SHINTANI, D. AND DELLAPENNA, D.: "A new paradigm for plant biochemistry: Nutritional Genomics. Vitamin E synthesis and human nutrition as an example.", Abstract of the oral presentation at the annual meeting of The American Society of Plant Physiology, 28.06.1998 XP002135562 the whole document	
P,X	SHINTANI, D. AND DELLAPENNA, D.: "Elevating the vitamin E content of plants through metabolic engineering" SCIENCE, vol. 282, no. 5396, 11 December 1998 (1998-12-11), pages 2098-2100, XP000887122 the whole document	18
P,A	WO 99 04622 A (UNIV NEVADA) 4 February 1999 (1999-02-04)	

INTERNATIONAL SEARCH REPORT

Inter. application No.
PCT/US 99/28588

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1, 3-11, 13-25 (all partially)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: Invention 1: Claims 1, 3-11, 13-25 (all partially)

Corn gamma-tocopherol methyltransferase:

Polynucleotide encoding a polypeptide as in SEQ ID NO: 2 or 22, polypeptide as in SEQ ID NO:2 or 22, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:1 or 21. Subject-matter relating to corn gamma-tocopherol methyltransferase.

2. Claims: Invention 2: Claims 1, 3-11, 13-25 (all partially)

Rice gamma-tocopherol methyltransferase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:4, 6, 24 or 26, polypeptide as in SEQ ID NO:4, 6, 24 or 26, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:3, 5, 23 or 25. Subject-matter relating to rice gamma-tocopherol methyltransferase.

3. Claims: Invention 3: Claims 1, 3-11, 13-25 (all partially)

Soybean gamma-tocopherol methyltransferase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:8 or 28, polypeptide as in SEQ ID NO:8 or 28, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:7 or 27. Subject-matter relating to soybean gamma-tocopherol methyltransferase.

4. Claims: Invention 4: Claims 1, 3-11, 13-25 (all partially)

Wheat gamma-tocopherol methyltransferase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:8 or 28, polypeptide as in SEQ ID NO:10 or 30, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:10 or 30. Subject-matter relating to wheat gamma-tocopherol methyltransferase.

5. Claims: Invention 5: Claims 2-10, 12-19,
21-25 (all partially)

Rice 4-hydroxyphenylpyruvate dioxygenase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:12, 14 or 34, polypeptide as in SEQ ID NO:12, 14 or 34, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:11, 13 or 33. Subject-matter relating to rice 4-hydroxyphenylpyruvate dioxygenase.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

6. Claims: Invention 6: Claims 2-10, 12-19,
21-25 (all partially)

Soybean 4-hydroxyphenylpyruvate dioxygenase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:16 or 36, polypeptide as in SEQ ID NO:16 or 36, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:15 or 35. Subject-matter relating to soybean 4-hydroxyphenylpyruvate dioxygenase.

7. Claims: Invention 7: Claims 2-10, 12-19,
21-25 (all partially)

Veronica 4-hydroxyphenylpyruvate dioxygenase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:18, polypeptide as in SEQ ID NO:18, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:17. Subject-matter relating to veronica 4-hydroxyphenylpyruvate dioxygenase.

8. Claims: Invention 8: Claims 2-10, 12-19,
21-25 (all partially)

Wheat 4-hydroxyphenylpyruvate dioxygenase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:20 or 38, polypeptide as in SEQ ID NO:20 or 38, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:19 or 37. Subject-matter relating to wheat 4-hydroxyphenylpyruvate dioxygenase.

9. Claims: Invention 9: Claims 2-10, 12-19,
21-25 (all partially)

Catalpa 4-hydroxyphenylpyruvate dioxygenase:

Polynucleotide encoding a polypeptide as in SEQ ID NO:32, polypeptide as in SEQ ID NO:32, polynucleotide comprising a nucleotide sequence as in SEQ ID NO:31. Subject-matter relating to catalpa 4-hydroxyphenylpyruvate dioxygenase.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internal Application No

PCT/US 99/28588

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
W0 9904622 A	04-02-1999	AU 8506198 A EP 1009812 A	16-02-1999 21-06-2000



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

[illegible]

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE

PLANT VITAMIN E BIOSYNTHETIC ENZYMES

This application claims the benefit of U.S. Provisional Application No. 60/110,781, filed December 3, 1998.

5

FIELD OF THE INVENTION

This invention is in the field of plant molecular biology. More specifically, this invention pertains to nucleic acid fragments encoding vitamin E biosynthetic enzymes in plants and seeds.

BACKGROUND OF THE INVENTION

10 Vitamin E (alpha-tocopherol) is an essential element in the mammalian diet since mammals can not synthesize plastoquinones or tocopherols. The first step in the formation of plastoquinones and tocopherols in plants is the formation of homogениstate from 4-hydroxyphenylpyruvate, a reaction catalyzed by 4-hydroxyphenylpyruvate dioxygenase (EC 1.13.11.27). Genetic mutants of this enzyme in *Arabidopsis* are deficient in both
15 carotenoids and tocopherols (Norris et al. (1995) *Plant Cell* 7:2139-2149). Carotenoids (xanthophylls) in corn endosperm are valuable coloring agents in poultry feeds; tocopherols are antioxidants which may be important in oil stability and human health. Generally present at low levels in plant tissues, 4-hydroxyphenylpyruvate dioxygenase catalyzes a complex, irreversible reaction. Only recently has 4-hydroxyphenylpyruvate dioxygenase
20 been purified to homogeneity from a plant source (Garcia et al. (1997 *Biochem. J.* 325:761-769). In plants, cDNAs encoding 4-hydroxyphenylpyruvate dioxygenase have been identified in carrots, barley and *Arabidopsis thaliana* with at least two different variants existing in this last plant. 4-Hydroxyphenylpyruvate dioxygenase is a known herbicide target (Mayonado et al. (1989) *Pestic. Biochem. Physiol.* 35:138-145; Schultz et al. (1993)
25 *FEBS lett.* 318:162-166; Secor (1994) *Plant Phys.* 106:1429-1433). Even though plastoquinones and tocopherols are not synthesized by mammals and bacteria 4-hydroxyphenylpyruvate dioxygenase activity is present, often at high levels, and involved in phenylalanine and tyrosine degradation. Among others, the cDNAs encoding 4-hydroxyphenylpyruvate dioxygenase have been identified in *Mycosphaerella graminicola*,
30 mice and *Coccidioides immitis*.

Plants synthesize alpha, beta, gamma and delta tocopherols of which alpha tocopherol (vitamin E) has the highest value for human nutrition. In soybean 7% of the tocopherols are vitamin E. Gamma tocopherol methyltransferase catalyzes the final step in vitamin E
35 synthesis and has been purified to homogeneity from pepper, marigold, *Euglena* and spinach. The gene encoding gamma tocopherol methyltransferase from *Synechocystis* was identified by mutating an open reading frame encoding a methyltransferase located in the operon containing the 4-hydroxyphenylpyruvate dioxygenase gene in the *Synechocystis*

genomic sequence (Shintani and Della Penna (1998) Abstract for the American Society of Plant Physiologists meeting in Madison, WI).

Since mammals can not synthesize tocopherols, the enzymes described here may be used for the discovery of new herbicides.

5

SUMMARY OF THE INVENTION

The present invention relates to isolated polynucleotides comprising a nucleotide sequence encoding a polypeptide of at least 160 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of a corn gamma-tocopherol methyltransferase polypeptide of SEQ ID
10 NOs:2 and 22, a rice gamma-tocopherol methyltransferase polypeptide of SEQ ID NOs:4, 6, 24, and 26, a soybean gamma-tocopherol methyltransferase polypeptide of SEQ ID NOs:8 and 28, and a wheat gamma-tocopherol methyltransferase polypeptide of SEQ ID NOs:10 and 30. The present invention also relates to an isolated polynucleotide comprising the complement of the nucleotide sequences described above.

15

The present invention relates to isolated polynucleotides comprising a nucleotide sequence encoding a polypeptide of at least 150 amino acids that has at least 95% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of a catalpa 4-hydroxyphenylpyruvate dioxygenase polypeptide of SEQ ID
20 NO:32, a rice 4-hydroxyphenylpyruvate dioxygenase polypeptide of SEQ ID NOs:12, 14, and 34, a soybean 4-hydroxyphenylpyruvate dioxygenase polypeptide of SEQ ID NOs:16 and 36, a Vernonia 4-hydroxyphenylpyruvate dioxygenase of SEQ ID NO:18, and a wheat 4-hydroxyphenylpyruvate dioxygenase polypeptide of SEQ ID NOs:20 and 38. The present invention also relates to an isolated polynucleotide comprising the complement of the nucleotide sequences described above.

25

It is preferred that the isolated polynucleotides of the claimed invention consist of a nucleic acid sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, and 37 that codes for the polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, and 38. The present invention also relates to an isolated polynucleotide
30 comprising a nucleotide sequences of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, and the complement of such nucleotide sequences.

35

The present invention relates to a chimeric gene comprising an isolated polynucleotide of the present invention operably linked to suitable regulatory sequences.

The present invention relates to an isolated host cell comprising a chimeric gene of the present invention or an isolated polynucleotide of the present invention. The host cell may be eukaryotic, such as a yeast or a plant cell, or prokaryotic, such as a bacterial cell. The

present invention also relates to a virus, preferably a baculovirus, comprising an isolated polynucleotide of the present invention or a chimeric gene of the present invention.

The present invention relates to a process for producing an isolated host cell comprising a chimeric gene of the present invention or an isolated polynucleotide of the present invention, the process comprising either transforming or transfecting an isolated
5 compatible host cell with a chimeric gene or isolated polynucleotide of the present invention.

The present invention relates to a gamma-tocopherol methyltransferase polypeptide of at least 160 amino acids comprising at least 80% homology based on the Clustal method of alignment compared to a polypeptide selected from the group consisting of SEQ ID NOs:2,
10 4, 6, 8, 10, 22, 24, 26, 28, and 30.

The present invention relates to a 4-hydroxyphenylpyruvate dioxygenase polypeptide of at least 150 amino acids comprising at least 95% homology based on the Clustal method of alignment compared to a polypeptide selected from the group consisting of SEQ ID NOs:12, 14, 16, 18, 20, 32, 34, 36, and 38.

15 The present invention relates to a method of selecting an isolated polynucleotide that affects the level of expression of a gamma-tocopherol methyltransferase or a 4-hydroxy-phenylpyruvate dioxygenase polypeptide in a host cell, preferably a plant cell, the method comprising the steps of: (a) constructing an isolated polynucleotide of the present invention or an isolated chimeric gene of the present invention; (b) introducing the isolated
20 polynucleotide or the isolated chimeric gene into a host cell; (c) measuring the level a gamma-tocopherol methyltransferase or a 4-hydroxyphenylpyruvate dioxygenase polypeptide in the host cell containing the isolated polynucleotide; and (d) comparing the level of a gamma-tocopherol methyltransferase or a 4-hydroxyphenylpyruvate dioxygenase polypeptide in the host cell containing the isolated polynucleotide with the level of a gamma-
25 tocopherol methyltransferase or a 4-hydroxyphenylpyruvate dioxygenase polypeptide in the host cell that does not contain the isolated polynucleotide.

The present invention relates to a method of obtaining a nucleic acid fragment encoding a substantial portion of a gamma-tocopherol methyltransferase or a 4-hydroxy-phenylpyruvate dioxygenase polypeptide gene, preferably a plant gamma-tocopherol
30 methyltransferase or 4-hydroxyphenylpyruvate dioxygenase polypeptide gene, comprising the steps of : synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, and the
35 complement of such nucleotide sequences; and amplifying a nucleic acid fragment (preferably a cDNA inserted in a cloning vector) using the oligonucleotide primer. The amplified nucleic acid fragment preferably will encode a portion of a gamma-tocopherol methyltransferase or a 4-hydroxyphenylpyruvate dioxygenase amino acid sequence.

The present invention also relates to a method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a 4-hydroxyphenylpyruvate dioxygenase or a gamma tocopherol methyltransferase polypeptide comprising the steps of: probing a cDNA or genomic library with an isolated polynucleotide of the present invention; identifying a DNA clone that hybridizes with an isolated polynucleotide of the present invention; isolating the identified DNA clone; and sequencing the cDNA or genomic fragment that comprises the isolated DNA clone.

A further embodiment of the instant invention is a method for evaluating at least one compound for its ability to inhibit the activity of a 4-hydroxyphenylpyruvate dioxygenase or a gamma tocopherol methyltransferase, the method comprising the steps of:

- (a) transforming a host cell with a chimeric gene comprising a nucleic acid fragment encoding a 4-hydroxyphenylpyruvate dioxygenase or a gamma tocopherol methyltransferase, operably linked to suitable regulatory sequences; (b) growing the transformed host cell under conditions that are suitable for expression of the chimeric gene wherein expression of the chimeric gene results in production of 4-hydroxyphenylpyruvate dioxygenase or gamma tocopherol methyltransferase in the transformed host cell;
- (c) optionally purifying the 4-hydroxyphenylpyruvate dioxygenase or the gamma tocopherol methyltransferase expressed by the transformed host cell; (d) treating the 4-hydroxyphenylpyruvate dioxygenase or the gamma tocopherol methyltransferase with a compound to be tested; and (e) comparing the activity of the 4-hydroxyphenylpyruvate dioxygenase or the gamma tocopherol methyltransferase that has been treated with a test compound to the activity of an untreated 4-hydroxyphenylpyruvate dioxygenase or gamma tocopherol methyltransferase, thereby selecting compounds with potential for inhibitory activity.

The present invention relates to a composition, such as a hybridization mixture, comprising an isolated polynucleotide of the present invention.

The present invention relates to an isolated polynucleotide of the present invention comprising at least one of 30 contiguous nucleotides derived from a nucleic acid sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, and 37.

The present invention relates to an expression cassette comprising an isolated polynucleotide of the present invention operably linked to a promoter.

The present invention relates to a method for positive selection of a transformed cell comprising: (a) transforming a host cell with the chimeric gene of the present invention or an expression cassette of the present invention; and (b) growing the transformed host cell, preferably plant cell, such as a monocot or a dicot, under conditions which allow expression of the gamma-tocopherol methyltransferase or a 4-hydroxyphenylpyruvate dioxygenase polynucleotide in an amount sufficient to complement a null mutant to provide a positive selection means.

BRIEF DESCRIPTION OF THE
DRAWINGS AND SEQUENCE DESCRIPTIONS

The invention can be more fully understood from the following detailed description and the accompanying drawings and Sequence Listing which form a part of this application.

5 Figure 1 shows a comparison of the amino acid sequences of the gamma-tocopherol methyltransferase from soybean contig assembled from clones sah1c.pk001.k8, sgc5c.pk001.m23, and sah1c.pk004.g2 (SEQ ID NO:8), corn contig assembled from clones cr1n.pk0179.f10:fis, cs1.pk0065.f2, cta1n.pk0031.d2, p0060.coran49r, and p0103.ciaay86r (SEQ ID NO:22, the 3'-end sequence of rice clone rls72.pk0010.g3 (SEQ ID NO:24), the 5'-
10 end sequence of clone rls72.pk0010.g3 (SEQ ID NO:26), soybean clone sah1c.pk004.g2 (SEQ ID NO:28), wheat clone wr1.pk0077.f1:fis (SEQ ID NO:30), *Synechocystis sp.* (NCBI General Identifier No. 1001725; SEQ ID NO:39), and *Arabidopsis thaliana* (NCBI General Identifier No. 4106538; SEQ ID NO:40). Dashes are used by the program to maximize the alignment.

15 Figure 2 shows a comparison of the amino acid sequences of the 4-hydroxyphenyl-pyruvate dioxygenase from catalpa clone ncs.pk0012.g1:fis (SEQ ID NO:32), soybean clone sgc5c.pk001.j9:fis (SEQ ID NO:36), wheat clone wdk4c.pk006.m9:fis (SEQ ID NO:38), *Hordeum vulgare* having NCBI General Identifier No. 3334222 (SEQ ID NO:41), *Daucus carota* having NCBI General Identifier No. 3334219 (SEQ ID NO:42), and *Arabidopsis*
20 *thaliana* having NCBI General Identifier No. 3334223 (SEQ ID NO:43). Dashes are used by the program to maximize the alignment.

Figure 3 depicts the distribution of the percent of alpha- and gamma-tocopherol content of 25 transgenic soybean lines transformed with the gamma tocopherol methyltransferase sequence from clone sah1c.pk001.k8.

25 Table 1 lists the polypeptides that are described herein, the designation of the cDNA clones that comprise the nucleic acid fragments encoding polypeptides representing all or a substantial portion of these polypeptides, and the corresponding identifier (SEQ ID NO:) as used in the attached Sequence Listing. The sequence descriptions and Sequence Listing attached hereto comply with the rules governing nucleotide and/or amino acid sequence
30 disclosures in patent applications as set forth in 37 C.F.R. §1.821-1.825.

TABLE 1
Vitamin E Biosynthetic Enzymes

Protein	Clone Designation	SEQ ID NO:	
		(Nucleotide)	(Amino Acid)
Corn Gamma Tocopherol Methyltransferase	Contig of: cr1n.pk0179.f10 cs1.pk0065.e1 cs1.pk0065.f2 cta1n.pk0031.d2	1	2
Rice Gamma Tocopherol Methyltransferase	rl0n.pk085.e11	3	4
Rice Gamma Tocopherol Methyltransferase	Contig of: rl0n.pk099.d24 rls72.pk0010.g3	5	6
Soybean Gamma Tocopherol Methyltransferase	Contig of: sah1c.pk001.k8 sgc5c.pk001.m23 sah1c.pk004.g2	7	8
Wheat Gamma Tocopherol Methyltransferase	Contig of: wr1.pk0077.f1 wr1.pk177.b11 wle1n.pk0065.h9 wre1n.pk0111.d6	9	10
Rice 4-Hydroxyphenylpyruvate Dioxygenase	rlr12.pk0019.g5	11	12
Rice 4-Hydroxyphenylpyruvate Dioxygenase	rlr12.pk0025.e4	13	14
Soybean 4-Hydroxyphenylpyruvate Dioxygenase	Contig of: sgc5c.pk001.j9 sgs1c.pk002.a8 sfl1.pk126.n15 sgs1c.pk003.o5 sgc4c.pk001.p2	15	16
Vernonia 4-Hydroxyphenylpyruvate Dioxygenase	vs1.pk0015.b2	17	18
Wheat 4-Hydroxyphenylpyruvate Dioxygenase	Contig of: wdk4c.pk006.m9 wlmk1.pk0021.h10 wlmk1.pk0019.e2 wlm0.pk0035.d5 wlmk1.pk0013.g11	19	20
Corn Gamma Tocopherol Methyltransferase	Contig of: cr1n.pk0179.f10: fis cs1.pk0065.f2 cta1n.pk0031.d2 p0060.coran49r p0103.ciaay86r	21	22
Rice Gamma Tocopherol Methyltransferase	rls72.pk0010.g3-3'	23	24

Protein	Clone Designation	SEQ ID NO:	
		(Nucleotide)	(Amino Acid)
Rice Gamma Tocopherol Methyltransferase	rls72.pk0010.g3-5'	25	26
Soybean Gamma Tocopherol Methyltransferase	sah1c.pk004.g2	27	28
Wheat Gamma Tocopherol Methyltransferase	wr1.pk0077.fl: fis	29	30
Catalpa 4-Hydroxyphenyl-pyruvate Dioxygenase	ncs.pk0012.g1: fis	31	32
Rice 4-Hydroxyphenylpyruvate Dioxygenase	rlr12.pk0025.e4: fis	33	34
Soybean 4-Hydroxyphenyl-pyruvate Dioxygenase	sgc5c.pk001.j9: fis	35	36
Wheat 4-Hydroxyphenylpyruvate Dioxygenase	wdk4c.pk006.m9: fis	37	38

The Sequence Listing contains the one letter code for nucleotide sequence characters and the three letter codes for amino acids as defined in conformity with the IUPAC-IUBMB standards described in *Nucleic Acids Res.* 13:3021-3030 (1985) and in the *Biochemical J.* 219 (No. 2):345-373 (1984) which are herein incorporated by reference. The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

DETAILED DESCRIPTION OF THE INVENTION

In the context of this disclosure, a number of terms shall be utilized. As used herein, a “polynucleotide” is a nucleotide sequence such as a nucleic acid fragment. A polynucleotide may be a polymer of RNA or DNA that is single- or double-stranded, that optionally contains synthetic, non-natural or altered nucleotide bases. A polynucleotide in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA, synthetic DNA, or mixtures thereof. An isolated polynucleotide of the present invention may include at least one of 60 contiguous nucleotides, preferably at least one of 40 contiguous nucleotides, most preferably one of at least 30 contiguous nucleotides derived from SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, or the complement of such sequences.

As used herein, “contig” refers to a nucleotide sequence that is assembled from two or more constituent nucleotide sequences that share common or overlapping regions of sequence homology. For example, the nucleotide sequences of two or more nucleic acid fragments can be compared and aligned in order to identify common or overlapping sequences. Where common or overlapping sequences exist between two or more nucleic

acid fragments, the sequences (and thus their corresponding nucleic acid fragments) can be assembled into a single contiguous nucleotide sequence.

As used herein, "substantially similar" refers to nucleic acid fragments wherein changes in one or more nucleotide bases results in substitution of one or more amino acids, but do not affect the functional properties of the polypeptide encoded by the nucleotide sequence. "Substantially similar" also refers to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate alteration of gene expression by gene silencing through for example antisense or co-suppression technology. "Substantially similar" also refers to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotides that do not substantially affect the functional properties of the resulting transcript vis-à-vis the ability to mediate gene silencing or alteration of the functional properties of the resulting protein molecule. It is therefore understood that the invention encompasses more than the specific exemplary nucleotide or amino acid sequences and includes functional equivalents thereof.

Substantially similar nucleic acid fragments may be selected by screening nucleic acid fragments representing subfragments or modifications of the nucleic acid fragments of the instant invention, wherein one or more nucleotides are substituted, deleted and/or inserted, for their ability to affect the level of the polypeptide encoded by the unmodified nucleic acid fragment in a plant or plant cell. For example, a substantially similar nucleic acid fragment representing at least one of 30 contiguous nucleotides derived from the instant nucleic acid fragment can be constructed and introduced into a plant or plant cell. The level of the polypeptide encoded by the unmodified nucleic acid fragment present in a plant or plant cell exposed to the substantially similar nucleic fragment can then be compared to the level of the polypeptide in a plant or plant cell that is not exposed to the substantially similar nucleic acid fragment.

For example, it is well known in the art that antisense suppression and co-suppression of gene expression may be accomplished using nucleic acid fragments representing less than the entire coding region of a gene, and by nucleic acid fragments that do not share 100% sequence identity with the gene to be suppressed. Moreover, alterations in a nucleic acid fragment which result in the production of a chemically equivalent amino acid at a given site, but do not effect the functional properties of the encoded polypeptide, are well known in the art. Thus, a codon for the amino acid alanine, a hydrophobic amino acid, may be substituted by a codon encoding another less hydrophobic residue, such as glycine, or a more hydrophobic residue, such as valine, leucine, or isoleucine. Similarly, changes which result in substitution of one negatively charged residue for another, such as aspartic acid for glutamic acid, or one positively charged residue for another, such as lysine for arginine, can also be expected to produce a functionally equivalent product. Nucleotide changes which

result in alteration of the N-terminal and C-terminal portions of the polypeptide molecule would also not be expected to alter the activity of the polypeptide. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products. Consequently, an isolated polynucleotide

5 comprising a nucleotide sequence of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37 and the complement of such nucleotide sequences may be used in methods of selecting an isolated polynucleotide that affects the expression of a polypeptide

10 (vitamin E biosynthetic enzyme) in a host cell. A method of selecting an isolated polynucleotide that affects the level of expression of a polypeptide in a host cell (eukaryotic, such as plant or yeast, prokaryotic such as bacterial, or viral) may comprise the steps of: constructing an isolated polynucleotide of the present invention or an isolated chimeric gene of the present invention; introducing the isolated polynucleotide or the isolated chimeric

15 gene into a host cell; measuring the level a polypeptide in the host cell containing the isolated polynucleotide; and comparing the level of a polypeptide in the host cell containing the isolated polynucleotide with the level of a polypeptide in a host cell that does not contain the isolated polynucleotide.

Moreover, substantially similar nucleic acid fragments may also be characterized by

20 their ability to hybridize. Estimates of such homology are provided by either DNA-DNA or DNA-RNA hybridization under conditions of stringency as is well understood by those skilled in the art (Hames and Higgins, Eds. (1985) *Nucleic Acid Hybridisation*, IRL Press, Oxford, U.K.). Stringency conditions can be adjusted to screen for moderately similar fragments, such as homologous sequences from distantly related organisms, to highly similar

25 fragments, such as genes that duplicate functional enzymes from closely related organisms. Post-hybridization washes determine stringency conditions. One set of preferred conditions uses a series of washes starting with 6X SSC, 0.5% SDS at room temperature for 15 min, then repeated with 2X SSC, 0.5% SDS at 45°C for 30 min, and then repeated twice with 0.2X SSC, 0.5% SDS at 50°C for 30 min. A more preferred set of stringent conditions uses

30 higher temperatures in which the washes are identical to those above except for the temperature of the final two 30 min washes in 0.2X SSC, 0.5% SDS was increased to 60°C. Another preferred set of highly stringent conditions uses two final washes in 0.1X SSC, 0.1% SDS at 65°C.

Substantially similar nucleic acid fragments of the instant invention may also be

35 characterized by the percent identity of the amino acid sequences that they encode to the amino acid sequences disclosed herein, as determined by algorithms commonly employed by those skilled in this art. Suitable nucleic acid fragments (isolated polynucleotides of the present invention) encode polypeptides that are at least about 70% identical, preferably at

least about 80% identical to the amino acid sequences reported herein. Preferred nucleic acid fragments encode amino acid sequences that are at least about 85% identical to the amino acid sequences reported herein. More preferred nucleic acid fragments encode amino acid sequences that are at least about 90% identical to the amino acid sequences reported
5 herein. Most preferred are nucleic acid fragments that encode amino acid sequences that are at least about 95% identical to the amino acid sequences reported herein. Suitable nucleic acid fragments not only have the above homologies but typically encode a polypeptide having at least about 50 amino acids, preferably at least about 100 amino acids, more preferably at least about 150 amino acids, still more preferably at least about 200 amino
10 acids, and most preferably at least about 250 amino acids. Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH
15 PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5.

A "substantial portion" of an amino acid or nucleotide sequence comprises an amino acid or a nucleotide sequence that is sufficient to afford putative identification of the protein or gene that the amino acid or nucleotide sequence comprises. Amino acid and nucleotide
20 sequences can be evaluated either manually by one skilled in the art, or by using computer-based sequence comparison and identification tools that employ algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul et al. (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/). In general, a sequence of ten or more contiguous amino acids or thirty or more contiguous nucleotides is necessary in order to putatively
25 identify a polypeptide or nucleic acid sequence as homologous to a known protein or gene. Moreover, with respect to nucleotide sequences, gene-specific oligonucleotide probes comprising 30 or more contiguous nucleotides may be used in sequence-dependent methods of gene identification (e.g., Southern hybridization) and isolation (e.g., *in situ* hybridization of bacterial colonies or bacteriophage plaques). In addition, short oligonucleotides of 12 or
30 more nucleotides may be used as amplification primers in PCR in order to obtain a particular nucleic acid fragment comprising the primers. Accordingly, a "substantial portion" of a nucleotide sequence comprises a nucleotide sequence that will afford specific identification and/or isolation of a nucleic acid fragment comprising the sequence. The instant specification teaches amino acid and nucleotide sequences encoding polypeptides that
35 comprise one or more particular plant proteins. The skilled artisan, having the benefit of the sequences as reported herein, may now use all or a substantial portion of the disclosed sequences for purposes known to those skilled in this art. Accordingly, the instant invention

comprises the complete sequences as reported in the accompanying Sequence Listing, as well as substantial portions of those sequences as defined above.

“Codon degeneracy” refers to divergence in the genetic code permitting variation of the nucleotide sequence without effecting the amino acid sequence of an encoded polypeptide. Accordingly, the instant invention relates to any nucleic acid fragment comprising a nucleotide sequence that encodes all or a substantial portion of the amino acid sequences set forth herein. The skilled artisan is well aware of the “codon-bias” exhibited by a specific host cell in usage of nucleotide codons to specify a given amino acid. Therefore, when synthesizing a nucleic acid fragment for improved expression in a host cell, it is desirable to design the nucleic acid fragment such that its frequency of codon usage approaches the frequency of preferred codon usage of the host cell.

“Synthetic nucleic acid fragments” can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form larger nucleic acid fragments which may then be enzymatically assembled to construct the entire desired nucleic acid fragment. “Chemically synthesized”, as related to nucleic acid fragment, means that the component nucleotides were assembled *in vitro*. Manual chemical synthesis of nucleic acid fragments may be accomplished using well established procedures, or automated chemical synthesis can be performed using one of a number of commercially available machines. Accordingly, the nucleic acid fragments can be tailored for optimal gene expression based on optimization of nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan appreciates the likelihood of successful gene expression if codon usage is biased towards those codons favored by the host. Determination of preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

“Gene” refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. “Native gene” refers to a gene as found in nature with its own regulatory sequences. “Chimeric gene” refers any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. “Endogenous gene” refers to a native gene in its natural location in the genome of an organism. A “foreign” gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A “transgene” is a gene that has been introduced into the genome by a transformation procedure.

“Coding sequence” refers to a nucleotide sequence that codes for a specific amino acid sequence. “Regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

“Promoter” refers to a nucleotide sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. The promoter sequence consists of proximal and more distal upstream elements, the latter elements often referred to as enhancers. Accordingly, an “enhancer” is a nucleotide sequence which can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic nucleotide segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters which cause a nucleic acid fragment to be expressed in most cell types at most times are commonly referred to as “constitutive promoters”. New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro and Goldberg (1989) *Biochemistry of Plants* 15:1-82. It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, nucleic acid fragments of different lengths may have identical promoter activity.

The “translation leader sequence” refers to a nucleotide sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the primary transcript to mRNA, mRNA stability or translation efficiency. Examples of translation leader sequences have been described (Turner and Foster (1995) *Mol. Biotechnol.* 3:225-236).

The “3' non-coding sequences” refer to nucleotide sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al. (1989) *Plant Cell* 1:671-680.

“RNA transcript” refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary

copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from posttranscriptional processing of the primary transcript and is referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into polypeptide by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA refers to an RNA transcript that includes the mRNA and so can be translated into a polypeptide by the cell. "Antisense RNA" refers to an RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene (see U.S. Patent No. 5,107,065, incorporated herein by reference). The complementarity of an antisense RNA may be with any part of the specific nucleotide sequence, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. "Functional RNA" refers to sense RNA, antisense RNA, ribozyme RNA, or other RNA that may not be translated but yet has an effect on cellular processes.

The term "operably linked" refers to the association of two or more nucleic acid fragments on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

The term "expression", as used herein, refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from the nucleic acid fragment of the invention. Expression may also refer to translation of mRNA into a polypeptide. "Antisense inhibition" refers to the production of antisense RNA transcripts capable of suppressing the expression of the target protein. "Overexpression" refers to the production of a gene product in transgenic organisms that exceeds levels of production in normal or non-transformed organisms. "Co-suppression" refers to the production of sense RNA transcripts capable of suppressing the expression of identical or substantially similar foreign or endogenous genes (U.S. Patent No. 5,231,020, incorporated herein by reference).

"Altered levels" refers to the production of gene product(s) in transgenic organisms in amounts or proportions that differ from that of normal or non-transformed organisms.

"Mature" protein refers to a post-translationally processed polypeptide; i.e., one from which any pre- or propeptides present in the primary translation product have been removed. "Precursor" protein refers to the primary product of translation of mRNA; i.e., with pre- and propeptides still present. Pre- and propeptides may be but are not limited to intracellular localization signals.

A "chloroplast transit peptide" is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the chloroplast or other plastid types present in the cell in which the protein is made. "Chloroplast transit sequence" refers to a

nucleotide sequence that encodes a chloroplast transit peptide. A "signal peptide" is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the secretory system (Chrispeels (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53). If the protein is to be directed to a vacuole, a vacuolar targeting signal (*supra*) can further be added, or if to the endoplasmic reticulum, an endoplasmic reticulum retention signal (*supra*) may be added. If the protein is to be directed to the nucleus, any signal peptide present should be removed and instead a nuclear localization signal included (Raikhel (1992) *Plant Phys.* 100:1627-1632).

"Transformation" refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" organisms. Examples of methods of plant transformation include *Agrobacterium*-mediated transformation (De Blaere et al. (1987) *Meth. Enzymol.* 143:277) and particle-accelerated or "gene gun" transformation technology (Klein et al. (1987) *Nature (London)* 327:70-73; U.S. Patent No. 4,945,050, incorporated herein by reference).

Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described more fully in Sambrook et al. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 1989 (hereinafter "Maniatis").

Nucleic acid fragments encoding at least a portion of several vitamin E biosynthetic enzymes have been isolated and identified by comparison of random plant cDNA sequences to public databases containing nucleotide and protein sequences using the BLAST algorithms well known to those skilled in the art. The nucleic acid fragments of the instant invention may be used to isolate cDNAs and genes encoding homologous proteins from the same or other plant species. Isolation of homologous genes using sequence-dependent protocols is well known in the art. Examples of sequence-dependent protocols include, but are not limited to, methods of nucleic acid hybridization, and methods of DNA and RNA amplification as exemplified by various uses of nucleic acid amplification technologies (e.g., polymerase chain reaction, ligase chain reaction).

For example, genes encoding other 4-hydroxyphenylpyruvate dioxygenases or gamma tocopherol methyltransferases, either as cDNAs or genomic DNAs, could be isolated directly by using all or a portion of the instant nucleic acid fragments as DNA hybridization probes to screen libraries from any desired plant employing methodology well known to those skilled in the art. Specific oligonucleotide probes based upon the instant nucleic acid sequences can be designed and synthesized by methods known in the art (Maniatis). Moreover, the entire sequences can be used directly to synthesize DNA probes by methods known to the skilled artisan such as random primer DNA labeling, nick translation, or end-labeling techniques, or RNA probes using available *in vitro* transcription systems. In

In addition, specific primers can be designed and used to amplify a part or all of the instant sequences. The resulting amplification products can be labeled directly during amplification reactions or labeled after amplification reactions, and used as probes to isolate full length cDNA or genomic fragments under conditions of appropriate stringency.

- 5 In addition, two short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols to amplify longer nucleic acid fragments encoding homologous genes from DNA or RNA. The polymerase chain reaction may also be performed on a library of cloned nucleic acid fragments wherein the sequence of one primer is derived from the instant nucleic acid fragments, and the sequence of the other primer takes
- 10 advantage of the presence of the polyadenylic acid tracts to the 3' end of the mRNA precursor encoding plant genes. Alternatively, the second primer sequence may be based upon sequences derived from the cloning vector. For example, the skilled artisan can follow the RACE protocol (Frohman et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:8998-9002) to generate cDNAs by using PCR to amplify copies of the region between a single point in the
- 15 transcript and the 3' or 5' end. Primers oriented in the 3' and 5' directions can be designed from the instant sequences. Using commercially available 3' RACE or 5' RACE systems (BRL), specific 3' or 5' cDNA fragments can be isolated (Ohara et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:5673-5677; Loh et al. (1989) *Science* 243:217-220). Products generated by the 3' and 5' RACE procedures can be combined to generate full-length cDNAs (Frohman
- 20 and Martin (1989) *Techniques* 1:165). Consequently, a polynucleotide comprising a nucleotide sequence of at least one of 60 (preferably one of at least 40, most preferably one of at least 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37 and the complement of such nucleotide sequences may be used in such methods to
- 25 obtain a nucleic acid fragment encoding a substantial portion of an amino acid sequence of a polypeptide. The present invention relates to a method of obtaining a nucleic acid fragment encoding a substantial portion of a polypeptide of a gene (such as gamma tocopherol methyltransferase or 4-phenylpyruvate dioxygenase) preferably a substantial portion of a plant polypeptide of a gene, comprising the steps of: synthesizing an oligonucleotide primer
- 30 comprising a nucleotide sequence of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, and the complement of such nucleotide sequences; and amplifying a nucleic acid fragment (preferably a cDNA inserted in a cloning vector) using the
- 35 oligonucleotide primer. The amplified nucleic acid fragment preferably will encode a portion of a polypeptide (gamma tocopherol methyltransferase or 4-phenylpyruvate dioxygenase).

Availability of the instant nucleotide and deduced amino acid sequences facilitates immunological screening of cDNA expression libraries. Synthetic peptides representing portions of the instant amino acid sequences may be synthesized. These peptides can be used to immunize animals to produce polyclonal or monoclonal antibodies with specificity for peptides or proteins comprising the amino acid sequences. These antibodies can be then be used to screen cDNA expression libraries to isolate full-length cDNA clones of interest (Lerner (1984) *Adv. Immunol.* 36:1-34; Maniatis).

The nucleic acid fragments of the instant invention may be used to create transgenic plants in which the disclosed polypeptides are present at higher or lower levels than normal or in cell types or developmental stages in which they are not normally found. This would have the effect of altering the level of homogenistate in those cells. Overexpression of 4-hydroxyphenylpyruvate dioxygenase should result in a larger accumulation of homogenistate which may be used by gamma tocopherol methyltransferase to produce vitamin E. Since mammals can not synthesize tocopherols, the enzymes described herein may be used for the discovery of new herbicides.

Overexpression of the proteins of the instant invention may be accomplished by first constructing a chimeric gene in which the coding region is operably linked to a promoter capable of directing expression of a gene in the desired tissues at the desired stage of development. For reasons of convenience, the chimeric gene may comprise promoter sequences and translation leader sequences derived from the same genes. 3' Non-coding sequences encoding transcription termination signals may also be provided. The instant chimeric gene may also comprise one or more introns in order to facilitate gene expression.

Plasmid vectors comprising the instant chimeric gene can then be constructed. The choice of plasmid vector is dependent upon the method that will be used to transform host plants. The skilled artisan is well aware of the genetic elements that must be present on the plasmid vector in order to successfully transform, select and propagate host cells containing the chimeric gene. The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression (Jones et al. (1985) *EMBO J.* 4:2411-2418; De Almeida et al. (1989) *Mol. Gen. Genetics* 218:78-86), and thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of DNA, Northern analysis of mRNA expression, Western analysis of protein expression, or phenotypic analysis.

For some applications it may be useful to direct the instant polypeptides to different cellular compartments, or to facilitate its secretion from the cell. It is thus envisioned that the chimeric gene described above may be further supplemented by directing the coding sequence to encode the instant polypeptides with appropriate intracellular targeting sequences such as transit sequences (Keegstra (1989) *Cell* 56:247-253), signal sequences or

sequences encoding endoplasmic reticulum localization (Chrispeels (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53), or nuclear localization signals (Raikhel (1992) *Plant Phys.* 100:1627-1632) with or without removing targeting sequences that are already present. While the references cited give examples of each of these, the list is not exhaustive and more
5 targeting signals of use may be discovered in the future.

It may also be desirable to reduce or eliminate expression of genes encoding the instant polypeptides in plants for some applications. In order to accomplish this, a chimeric gene designed for co-suppression of the instant polypeptide can be constructed by linking a gene or gene fragment encoding that polypeptide to plant promoter sequences.
10 Alternatively, a chimeric gene designed to express antisense RNA for all or part of the instant nucleic acid fragment can be constructed by linking the gene or gene fragment in reverse orientation to plant promoter sequences. Either the co-suppression or antisense chimeric genes could be introduced into plants via transformation wherein expression of the corresponding endogenous genes are reduced or eliminated.

15 Molecular genetic solutions to the generation of plants with altered gene expression have a decided advantage over more traditional plant breeding approaches. Changes in plant phenotypes can be produced by specifically inhibiting expression of one or more genes by antisense inhibition or cosuppression (U.S. Patent Nos. 5,190,931, 5,107,065 and 5,283,323). An antisense or cosuppression construct would act as a dominant negative
20 regulator of gene activity. While conventional mutations can yield negative regulation of gene activity these effects are most likely recessive. The dominant negative regulation available with a transgenic approach may be advantageous from a breeding perspective. In addition, the ability to restrict the expression of specific phenotype to the reproductive tissues of the plant by the use of tissue specific promoters may confer agronomic advantages
25 relative to conventional mutations which may have an effect in all tissues in which a mutant gene is ordinarily expressed.

The person skilled in the art will know that special considerations are associated with the use of antisense or cosuppression technologies in order to reduce expression of particular genes. For example, the proper level of expression of sense or antisense genes may require
30 the use of different chimeric genes utilizing different regulatory elements known to the skilled artisan. Once transgenic plants are obtained by one of the methods described above, it will be necessary to screen individual transgenics for those that most effectively display the desired phenotype. Accordingly, the skilled artisan will develop methods for screening large numbers of transformants. The nature of these screens will generally be chosen on
35 practical grounds, and is not an inherent part of the invention. For example, one can screen by looking for changes in gene expression by using antibodies specific for the protein encoded by the gene being suppressed, or one could establish assays that specifically measure enzyme activity. A preferred method will be one which allows large numbers of

samples to be processed rapidly, since it will be expected that a large number of transformants will be negative for the desired phenotype.

The instant polypeptides (or portions thereof) may be produced in heterologous host cells, particularly in the cells of microbial hosts, and can be used to prepare antibodies to these proteins by methods well known to those skilled in the art. The antibodies are useful for detecting the polypeptides of the instant invention *in situ* in cells or *in vitro* in cell extracts. Preferred heterologous host cells for production of the instant polypeptides are microbial hosts. Microbial expression systems and expression vectors containing regulatory sequences that direct high level expression of foreign proteins are well known to those skilled in the art. Any of these could be used to construct a chimeric gene for production of the instant polypeptides. This chimeric gene could then be introduced into appropriate microorganisms via transformation to provide high level expression of the encoded vitamin E biosynthetic enzyme. An example of a vector for high level expression of the instant polypeptides in a bacterial host is provided (Example 7).

Additionally, the instant polypeptides can be used as a targets to facilitate design and/or identification of inhibitors of those enzymes that may be useful as herbicides. This is desirable because the polypeptides described herein catalyze various steps in vitamin E biosynthesis. Accordingly, inhibition of the activity of one or more of the enzymes described herein could lead to inhibition of plant growth. Thus, the instant polypeptides could be appropriate for new herbicide discovery and design.

All or a substantial portion of the nucleic acid fragments of the instant invention may also be used as probes for genetically and physically mapping the genes that they are a part of, and as markers for traits linked to those genes. Such information may be useful in plant breeding in order to develop lines with desired phenotypes. For example, the instant nucleic acid fragments may be used as restriction fragment length polymorphism (RFLP) markers. Southern blots (Maniatis) of restriction-digested plant genomic DNA may be probed with the nucleic acid fragments of the instant invention. The resulting banding patterns may then be subjected to genetic analyses using computer programs such as MapMaker (Lander et al. (1987) *Genomics* 1:174-181) in order to construct a genetic map. In addition, the nucleic acid fragments of the instant invention may be used to probe Southern blots containing restriction endonuclease-treated genomic DNAs of a set of individuals representing parent and progeny of a defined genetic cross. Segregation of the DNA polymorphisms is noted and used to calculate the position of the instant nucleic acid sequence in the genetic map previously obtained using this population (Botstein et al. (1980) *Am. J. Hum. Genet.* 32:314-331).

The production and use of plant gene-derived probes for use in genetic mapping is described in Bernatzky and Tanksley (1986) *Plant Mol. Biol. Reporter* 4:37-41. Numerous publications describe genetic mapping of specific cDNA clones using the methodology

outlined above or variations thereof. For example, F2 intercross populations, backcross populations, randomly mated populations, near isogenic lines, and other sets of individuals may be used for mapping. Such methodologies are well known to those skilled in the art.

5 Nucleic acid probes derived from the instant nucleic acid sequences may also be used for physical mapping (i.e., placement of sequences on physical maps; see Hoheisel et al. In: *Nonmammalian Genomic Analysis: A Practical Guide*, Academic press 1996, pp. 319-346, and references cited therein).

10 In another embodiment, nucleic acid probes derived from the instant nucleic acid sequences may be used in direct fluorescence *in situ* hybridization (FISH) mapping (Trask (1991) *Trends Genet.* 7:149-154). Although current methods of FISH mapping favor use of large clones (several to several hundred KB; see Laan et al. (1995) *Genome Res.* 5:13-20), improvements in sensitivity may allow performance of FISH mapping using shorter probes.

A variety of nucleic acid amplification-based methods of genetic and physical mapping may be carried out using the instant nucleic acid sequences. Examples include
15 allele-specific amplification (Kazazian (1989) *J. Lab. Clin. Med.* 11:95-96), polymorphism of PCR-amplified fragments (CAPS; Sheffield et al. (1993) *Genomics* 16:325-332), allele-specific ligation (Landegren et al. (1988) *Science* 241:1077-1080), nucleotide extension reactions (Sokolov (1990) *Nucleic Acid Res.* 18:3671), Radiation Hybrid Mapping (Walter et al. (1997) *Nat. Genet.* 7:22-28) and Happy Mapping (Dear and Cook (1989) *Nucleic Acid*
20 *Res.* 17:6795-6807). For these methods, the sequence of a nucleic acid fragment is used to design and produce primer pairs for use in the amplification reaction or in primer extension reactions. The design of such primers is well known to those skilled in the art. In methods employing PCR-based genetic mapping, it may be necessary to identify DNA sequence differences between the parents of the mapping cross in the region corresponding to the
25 instant nucleic acid sequence. This, however, is generally not necessary for mapping methods.

Loss of function mutant phenotypes may be identified for the instant cDNA clones either by targeted gene disruption protocols or by identifying specific mutants for these genes contained in a maize population carrying mutations in all possible genes (Ballinger
30 and Benzer (1989) *Proc. Natl. Acad. Sci USA* 86:9402-9406; Koes et al. (1995) *Proc. Natl. Acad. Sci USA* 92:8149-8153; Bensen et al. (1995) *Plant Cell* 7:75-84). The latter approach may be accomplished in two ways. First, short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols in conjunction with a mutation tag sequence primer on DNAs prepared from a population of plants in which
35 Mutator transposons or some other mutation-causing DNA element has been introduced (see Bensen, *supra*). The amplification of a specific DNA fragment with these primers indicates the insertion of the mutation tag element in or near the plant gene encoding the instant polypeptides. Alternatively, the instant nucleic acid fragment may be used as a

- hybridization probe against PCR amplification products generated from the mutation population using the mutation tag sequence primer in conjunction with an arbitrary genomic site primer, such as that for a restriction enzyme site-anchored synthetic adaptor. With either method, a plant containing a mutation in the endogenous gene encoding the instant polypeptides can be identified and obtained. This mutant plant can then be used to determine or confirm the natural function of the instant polypeptides disclosed herein.

EXAMPLES

- The present invention is further defined in the following Examples, in which all parts and percentages are by weight and degrees are Celsius, unless otherwise stated. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

EXAMPLE 1

Composition of cDNA Libraries; Isolation and Sequencing of cDNA Clones

cDNA libraries representing mRNAs from various corn, rice, soybean, *Vernonia*, and wheat tissues were prepared. The characteristics of the libraries are described below.

TABLE 2

cDNA Libraries from cDNA Libraries from Corn, Rice, Soybean, *Vernonia*, and Wheat

Library	Tissue	Clone
cr1n	Corn Root From 7 Day Old Seedlings*	cr1n.pk0155.d1
cr1n	Corn Root From 7 Day Old Seedlings*	cr1n.pk0179.f10
cs1	Corn Leaf Sheath From 5 Week Old Plant	cs1.pk0065.e1
cs1	Corn Leaf Sheath From 5 Week Old Plant	cs1.pk0065.f2
cta1n	Corn Tassel*	cta1n.pk0031.d2
ncs	<i>Catalpa speciosa</i> Developing Seed	ncs.pk0012.g1
p0023	Corn Leaf From Plant Transformed with Gene MIC07 (leucine-rich repeat) Which Induces Resistance Prior to Genetic Lesion Formation. Harvested About One Month After Planting in Green House*	p0060.coran49r
p0103	Corn Tassel Shoots(0.1-1.4 cm)*	p0103.ciaay86r
rl0n	Rice 15 Day Old Leaf*	rl0n.pk085.e11
rl0n	Rice 15 Day Old Leaf*	rl0n.pk099.d24
rlr12	Rice Leaf 15 Days After Germination, 12 Hours After Infection of Strain <i>Magaporthe grisea</i> 4360-R-62 (AVR2-YAMO); Resistant	rlr12.pk0019.g5

Library	Tissue	Clone
rlr12	Rice Leaf 15 Days After Germination, 12 Hours After Infection of Strain <i>Magaporthe grisea</i> 4360-R-62 (AVR2-YAMO); Resistant	rlr12.pk0025.e4
rls72	Rice Leaf 15 Days After Germination, 72 Hours After Infection of Strain <i>Magaporthe grisea</i> 4360-R-67 (AVR2-YAMO); Susceptible	rls72.pk0010.g3
sah1c	Soybean Sprayed With Authority Herbicide	sah1c.pk001.k8
sah1c	Soybean Sprayed With Authority Herbicide	sah1c.pk004.g2
sfl1	Soybean Immature Flower	sfl1.pk126.n15
sgc4c	Soybean Cotyledon 14-21 Days After Germination (1/4 yellow)	sgc4c.pk001.p2
sgc5c	Soybean (Cotyledon 15-24 Days After Germination (3/4 yellow)	sgc5c.pk001.j9
sgc5c	Soybean (Cotyledon 15-24 Days After Germination (3/4 yellow)	sgc5c.pk001.m23
sgs1c	Soybean Seeds 4 Hours After Germination	sgs1c.pk002.a8
sgs1c	Soybean Seeds 4 Hours After Germination	sgs1c.pk003.o5
vs1	<i>Vernonia</i> Seed Stage 1	vs1.pk0015.b2
wdk4c	Wheat Developing Kernel, 21 Days After Anthesis	wdk4c.pk006.m9
wle1n	Wheat Leaf From 7 Day Old Etiolated Seedling*	wle1n.pk0065.h9
wlm0	Wheat Seedlings 0 Hour After Inoculation With <i>Erysiphe graminis f. sp tritici</i>	wlm0.pk0035.d5
wlmk1	Wheat Seedlings 1 Hour After Inoculation With <i>Erysiphe graminis f. sp tritici</i> and Treatment With Herbicide**	wlmk1.pk0013.g11
wlmk1	Wheat Seedlings 1 Hour After Inoculation With <i>Erysiphe graminis f. sp tritici</i> and Treatment With Herbicide**	wlmk1.pk0019.e2
wlmk1	Wheat Seedlings 1 Hour After Inoculation With <i>Erysiphe graminis f. sp tritici</i> and Treatment With Herbicide**	wlmk1.pk0021.h10
wr1	Wheat Root From 7 Day Old Seedling	wr1.pk0077.f1
wr1	Wheat Root From 7 Day Old Seedling	wr1.pk177.b11
wre1n	Wheat Root From 7 Day Old Etiolated Seedling*	wre1n.pk0111.d6

* These libraries were normalized essentially as described in U.S. Patent No. 5,482,845, incorporated herein by reference.

5 ** Application of 6-iodo-2-propoxy-3-propyl-4(3H)-quinazolinone; synthesis and methods of using this compound are described in USSN 08/545,827, incorporated herein by reference.

10 cDNA libraries may be prepared by any one of many methods available. For example, the cDNAs may be introduced into plasmid vectors by first preparing the cDNA libraries in Uni-ZAP™ XR vectors according to the manufacturer's protocol (Stratagene

Cloning Systems, La Jolla, CA). The Uni-ZAP™ XR libraries are converted into plasmid libraries according to the protocol provided by Stratagene. Upon conversion, cDNA inserts will be contained in the plasmid vector pBluescript. In addition, the cDNAs may be introduced directly into precut Bluescript II SK(+) vectors (Stratagene) using T4 DNA ligase (New England Biolabs), followed by transfection into DH10B cells according to the manufacturer's protocol (GIBCO BRL Products). Once the cDNA inserts are in plasmid vectors, plasmid DNAs are prepared from randomly picked bacterial colonies containing recombinant pBluescript plasmids, or the insert cDNA sequences are amplified via polymerase chain reaction using primers specific for vector sequences flanking the inserted cDNA sequences. Amplified insert DNAs or plasmid DNAs are sequenced in dye-primer sequencing reactions to generate partial cDNA sequences (expressed sequence tags or "ESTs"; see Adams et al., (1991) *Science* 252:1651-1656). The resulting ESTs are analyzed using a Perkin Elmer Model 377 fluorescent sequencer.

EXAMPLE 2

Identification of cDNA Clones

cDNA clones encoding vitamin E biosynthetic enzymes were identified by conducting BLAST (Basic Local Alignment Search Tool; Altschul et al. (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/) searches for similarity to sequences contained in the BLAST "nr" database (comprising all non-redundant GenBank CDS translations, sequences derived from the 3-dimensional structure Brookhaven Protein Data Bank, the last major release of the SWISS-PROT protein sequence database, EMBL, and DDBJ databases). The cDNA sequences obtained in Example 1 were analyzed for similarity to all publicly available DNA sequences contained in the "nr" database using the BLASTN algorithm provided by the National Center for Biotechnology Information (NCBI). The DNA sequences were translated in all reading frames and compared for similarity to all publicly available protein sequences contained in the "nr" database using the BLASTX algorithm (Gish and States (1993) *Nat. Genet.* 3:266-272) provided by the NCBI. For convenience, the P-value (probability) of observing a match of a cDNA sequence to a sequence contained in the searched databases merely by chance as calculated by BLAST are reported herein as "pLog" values, which represent the negative of the logarithm of the reported P-value. Accordingly, the greater the pLog value, the greater the likelihood that the cDNA sequence and the BLAST "hit" represent homologous proteins.

EXAMPLE 3

Characterization of cDNA Clones Encoding

Gamma Tocopherol Methyltransferase

The BLASTX search using the EST sequences from clones listed in Table 3 revealed similarity of the polypeptides encoded by the cDNAs to gamma tocopherol methyltransferase from *Synechocystis* sp. (NCBI General Identifier No. 1001725). Shown in

Table 3 are the BLAST results for individual ESTs ("EST"), the sequences of the entire cDNA inserts comprising the indicated cDNA clones ("FIS"), contigs assembled from two or more ESTs ("Contig"), contigs assembled from an FIS and one or more ESTs ("Contig*"), or sequences encoding the entire protein derived from an FIS, a contig, or an FIS and PCR ("CGS"):

TABLE 3

BLAST Results for Sequences Encoding Polypeptides Homologous to gamma Tocopherol Methyltransferase

Clone	Status	BLAST pLog Score 1001725
Contig of: cr1n.pk0179.f10 cs1.pk0065.e1 cs1.pk0065.f2 cta1n.pk0031.d2	Contig	49.70
rl0n.pk085.e11	EST	15.10
Contig of: rl0n.pk099.d24 rls72.pk0010.g3	Contig	8.30
Contig of: sah1c.pk001.k8 sgc5c.pk001.m23 sah1c.pk004.g2:fis	CGS	64.22
Contig of: wr1.pk0077.f1 wr1.pk177.b11 wle1n.pk0065.h9 wre1n.pk0111.d6	Contig	57.00

10

Further sequencing and searching of the DuPont proprietary database allowed for the assembly of longer sequences. The BLASTX search using the nucleotide sequences from clones listed in Table 4 revealed similarity of the polypeptides encoded by the cDNAs to gamma tocopherol methyltransferase from *Arabidopsis thaliana* (NCBI General Identifier No. 4106538). Shown in Table 4 are the BLAST results for individual ESTs ("EST") or sequences encoding the entire protein derived from the sequences of the entire cDNA inserts comprising the indicated cDNA clones ("FIS"), contigs assembled from an FIS and one or more ESTs, or an FIS and PCR ("CGS"):

15

TABLE 4

BLAST Results for Sequences Encoding Polypeptides Homologous
to gamma Tocopherol Methyltransferase

Clone	Status	BLAST pLog Score 4106538
Contig of: cr1n.pk0179.f10: fis cs1.pk0065.f2 cta1n.pk0031.d2 p0060.coran49r p0103.ciaay86r	CGS	118.00
rls72.pk0010.g3-3'	EST*	81.30
rls72.pk0010.g3-5'	EST*	11.00
sah1c.pk004.g2	CGS	125.00
wr1.pk0077.f1: fis	CGS	120.00

*These sequences are derived from partial full-insert sequencing of clone rls72.pk0010.g3
and correspond to 5'-end and 3'-end sequences.

Figure 1 presents an alignment of the amino acid sequences set forth in SEQ ID
NOs:8, 22, 24, 26, 28, and 30 and the *Synechocystis sp.* and *Arabidopsis thaliana* sequences
(SEQ ID NO:39 and SEQ ID NO:40, respectively). The data in Table 5 represents a
calculation of the percent identity of the amino acid sequences set forth in SEQ ID NOs:8,
22, 24, 26, 28, and 30 and the *Synechocystis sp.* and *Arabidopsis thaliana* sequences (SEQ
ID NO:39 and SEQ ID NO:40, respectively).

TABLE 5

Percent Identity of Amino Acid Sequences Deduced From the Nucleotide
Sequences of cDNA Clones Encoding Polypeptides Homologous
to gamma Tocopherol Methyltransferase

SEQ ID NO.	Percent Identity to	
	1001725	4106538
8	37.9	56.9
22	42.6	58.6
24	42.8	67.6
26	23.8	33.3
28	41.3	62.6
30	43.2	59.5

Sequence alignments and percent identity calculations were performed using the
Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc.,
Madison, WI). Multiple alignment of the sequences was performed using the Clustal
method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default

- parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. Sequence alignments and BLAST scores and probabilities indicate that the nucleic acid fragments comprising the instant cDNA clones
- 5 encode one entire corn, two entire soybean, one entire wheat, and two fragments corresponding to an almost entire rice gamma tocopherol methyltransferase. These sequences represent the first corn, rice, soybean, and wheat sequences encoding gamma tocopherol methyltransferase.

EXAMPLE 4

10 Characterization of cDNA Clones Encoding 4-Hydroxyphenylpyruvate Dioxygenase

- The BLASTX search using the EST sequences from clones listed in Table 6 revealed similarity of the polypeptides encoded by the cDNAs to 4-hydroxyphenylpyruvate dioxygenase from *Hordeum vulgare* (NCBI General Identifier No. 3334222), *Daucus carota*
- 15 (NCBI General Identifier No. 3334219) and *Arabidopsis thaliana* (NCBI General Identifier No. 3334223). Shown in Table 6 are the BLAST results for individual ESTs ("EST") or contigs assembled from two or more ESTs ("Contig"):

TABLE 6

20 BLAST Results for Sequences Encoding Polypeptides Homologous
to 4-Hydroxyphenylpyruvate Dioxygenase

Clone	Status	BLAST pLog Score		
		3334222	3334219	3334223
rlr12.pk0019.g5	EST	26.70	17.70	18.00
rlr12.pk0025.e4	EST	58.52	50.40	49.40
Contig of: sgc5c.pk001.j9 sgs1c.pk002.a8 sfl1.pk126.n15 sgs1c.pk003.o5 sgc4c.pk001.p2	Contig	55.00	81.30	107.00
vs1.pk0015.b2	EST	39.40	51.70	52.05
Contig of: wdk4c.pk006.m9 wlmk1.pk0021.h1	Contig	176.00	102.00	97.00
0 wlmk1.pk0019.e2 wlm0.pk0035.d5 wlmk1.pk0013.g1				
1				

Further sequencing and searching of the DuPont proprietary database allowed the assembly of longer sequences and identification of 4-hydroxyphenylpyruvate dioxygenase in

- other species. The BLASTX search using the nucleotide sequences from clones listed in Table 7 revealed similarity of the polypeptides encoded by the cDNAs to 4-hydroxyphenylpyruvate dioxygenase from *Hordeum vulgare* (NCBI General Identifier No. 3334222), *Daucus carota* (NCBI General Identifier No. 3334219) and *Arabidopsis thaliana* (NCBI General Identifier No. 3334223). Shown in Table 7 are the BLAST results for the sequences of the entire cDNA inserts comprising the indicated cDNA clones ("FIS") or sequences encoding the entire protein derived from an FIS, or an FIS and PCR ("CGS"):

TABLE 7

BLAST Results for Sequences Encoding Polypeptides Homologous to 4-Hydroxyphenylpyruvate Dioxygenase

Clone	Status	BLAST pLog Score		
		3334222	3334219	3334223
ncs.pk0012.g1: fis	CGS	147.00	254.00	177.00
rlr12.pk0025.e4: fis	FIS	117.00	102.00	100.00
sgc5c.pk001.j9: fis	CGS	149.00	>254.00	>254.00
wdk4c.pk006.m9: fis	CGS	>254.00	155.00	151.00

- Figure 2 presents an alignment of the amino acid sequences set forth in SEQ ID NOs:32, 34, 36, and 38 and the *Hordeum vulgare*, *Daucus carota*, and *Arabidopsis thaliana* sequences (SEQ ID NO:41, SEQ ID NO:42, and SEQ ID NO:43, respectively). The data in Table 8 represents a calculation of the percent identity of the amino acid sequences set forth in SEQ ID NOs:18, 32, 34, 36, and 38 and the *Hordeum vulgare*, *Daucus carota*, and *Arabidopsis thaliana* sequences (SEQ ID NO:41, SEQ ID NO:42, and SEQ ID NO:43, respectively)

TABLE 8

Percent Identity of Amino Acid Sequences Deduced From the Nucleotide Sequences of cDNA Clones Encoding Polypeptides Homologous to 4-Hydroxyphenylpyruvate Dioxygenase

SEQ ID NO.	Percent Identity to		
	3334222	3334219	3334223
18	49.4	58.2	60.0
32	58.5	73.3	67.4
34	86.4	73.3	69.9
36	49.4	58.2	60.0
38	94.2	59.9	58.1

Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc.,

Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. Sequence alignments and BLAST scores and probabilities indicate that the nucleic acid fragments comprising the instant cDNA clones encode substantial portions of rice, soybean, *Vernonia*, and wheat 4-hydroxyphenylpyruvate dioxygenase, and entire catalpa, soybean, and wheat 4-hydroxyphenylpyruvate dioxygenase. These sequences represent the first catalpa, rice, soybean, *Vernonia*, and wheat sequences encoding 4-hydroxyphenylpyruvate dioxygenase.

EXAMPLE 5

Expression of Chimeric Genes in Monocot Cells

A chimeric gene comprising a cDNA encoding the instant polypeptides in sense orientation with respect to the maize 27 kD zein promoter that is located 5' to the cDNA fragment, and the 10 kD zein 3' end that is located 3' to the cDNA fragment, can be constructed. The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites (NcoI or SmaI) can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the digested vector pML103 as described below. Amplification is then performed in a standard PCR. The amplified DNA is then digested with restriction enzymes NcoI and SmaI and fractionated on an agarose gel. The appropriate band can be isolated from the gel and combined with a 4.9 kb NcoI-SmaI fragment of the plasmid pML103. Plasmid pML103 has been deposited under the terms of the Budapest Treaty at ATCC (American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209), and bears accession number ATCC 97366. The DNA segment from pML103 contains a 1.05 kb Sall-NcoI promoter fragment of the maize 27 kD zein gene and a 0.96 kb SmaI-Sall fragment from the 3' end of the maize 10 kD zein gene in the vector pGem9Zf(+) (Promega). Vector and insert DNA can be ligated at 15°C overnight, essentially as described (Maniatis). The ligated DNA may then be used to transform *E. coli* XL1-Blue (Epicurian Coli XL-1 Blue™; Stratagene). Bacterial transformants can be screened by restriction enzyme digestion of plasmid DNA and limited nucleotide sequence analysis using the dideoxy chain termination method (Sequenase™ DNA Sequencing Kit; U.S. Biochemical). The resulting plasmid construct would comprise a chimeric gene encoding, in the 5' to 3' direction, the maize 27 kD zein promoter, a cDNA fragment encoding the instant polypeptides, and the 10 kD zein 3' region.

The chimeric gene described above can then be introduced into corn cells by the following procedure. Immature corn embryos can be dissected from developing caryopses derived from crosses of the inbred corn lines H99 and LH132. The embryos are isolated

to 11 days after pollination when they are 1.0 to 1.5 mm long. The embryos are then placed with the axis-side facing down and in contact with agarose-solidified N6 medium (Chu et al. (1975) *Sci. Sin. Peking* 18:659-668). The embryos are kept in the dark at 27°C. Friable embryogenic callus consisting of undifferentiated masses of cells with somatic

- 5 proembryoids and embryoids borne on suspensor structures proliferates from the scutellum of these immature embryos. The embryogenic callus isolated from the primary explant can be cultured on N6 medium and sub-cultured on this medium every 2 to 3 weeks.

The plasmid, p35S/Ac (obtained from Dr. Peter Eckes, Hoechst Ag, Frankfurt, Germany) may be used in transformation experiments in order to provide for a selectable
10 marker. This plasmid contains the *Pat* gene (see European Patent Publication 0 242 236) which encodes phosphinothricin acetyl transferase (PAT). The enzyme PAT confers resistance to herbicidal glutamine synthetase inhibitors such as phosphinothricin. The *pat* gene in p35S/Ac is under the control of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812) and the 3' region of the nopaline synthase gene
15 from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*.

The particle bombardment method (Klein et al. (1987) *Nature* 327:70-73) may be used to transfer genes to the callus culture cells. According to this method, gold particles (1 µm in diameter) are coated with DNA using the following technique. Ten µg of plasmid DNAs are added to 50 µL of a suspension of gold particles (60 mg per mL). Calcium chloride
20 (50 µL of a 2.5 M solution) and spermidine free base (20 µL of a 1.0 M solution) are added to the particles. The suspension is vortexed during the addition of these solutions. After 10 minutes, the tubes are briefly centrifuged (5 sec at 15,000 rpm) and the supernatant removed. The particles are resuspended in 200 µL of absolute ethanol, centrifuged again and the supernatant removed. The ethanol rinse is performed again and the particles
25 resuspended in a final volume of 30 µL of ethanol. An aliquot (5 µL) of the DNA-coated gold particles can be placed in the center of a Kapton™ flying disc (Bio-Rad Labs). The particles are then accelerated into the corn tissue with a Biolistic™ PDS-1000/He (Bio-Rad Instruments, Hercules CA), using a helium pressure of 1000 psi, a gap distance of 0.5 cm and a flying distance of 1.0 cm.

30 For bombardment, the embryogenic tissue is placed on filter paper over agarose-solidified N6 medium. The tissue is arranged as a thin lawn and covered a circular area of about 5 cm in diameter. The petri dish containing the tissue can be placed in the chamber of the PDS-1000/He approximately 8 cm from the stopping screen. The air in the chamber is then evacuated to a vacuum of 28 inches of Hg. The macrocarrier is accelerated with a
35 helium shock wave using a rupture membrane that bursts when the He pressure in the shock tube reaches 1000 psi.

Seven days after bombardment the tissue can be transferred to N6 medium that contains gluphosinate (2 mg per liter) and lacks casein or proline. The tissue continues to

grow slowly on this medium. After an additional 2 weeks the tissue can be transferred to fresh N6 medium containing glufosinate. After 6 weeks, areas of about 1 cm in diameter of actively growing callus can be identified on some of the plates containing the glufosinate-supplemented medium. These calli may continue to grow when sub-cultured on the selective medium.

Plants can be regenerated from the transgenic callus by first transferring clusters of tissue to N6 medium supplemented with 0.2 mg per liter of 2,4-D. After two weeks the tissue can be transferred to regeneration medium (Fromm et al. (1990) *Bio/Technology* 8:833-839).

EXAMPLE 6

Expression of Chimeric Genes in Dicot Cells

A seed-specific expression cassette composed of the promoter and transcription terminator from the gene encoding the β subunit of the seed storage protein phaseolin from the bean *Phaseolus vulgaris* (Doyle et al. (1986) *J. Biol. Chem.* 261:9228-9238) can be used for expression of the instant polypeptides in transformed soybean. The phaseolin cassette includes about 500 nucleotides upstream (5') from the translation initiation codon and about 1650 nucleotides downstream (3') from the translation stop codon of phaseolin. Between the 5' and 3' regions are the unique restriction endonuclease sites Nco I (which includes the ATG translation initiation codon), Sma I, Kpn I and Xba I. The entire cassette is flanked by Hind III sites.

The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the expression vector. Amplification is then performed as described above, and the isolated fragment is inserted into a pUC18 vector carrying the seed expression cassette.

Soybean embryos may then be transformed with the expression vector comprising sequences encoding the instant polypeptides. To induce somatic embryos, cotyledons, 3-5 mm in length dissected from surface sterilized, immature seeds of the soybean cultivar A2872, can be cultured in the light or dark at 26°C on an appropriate agar medium for 6-10 weeks. Somatic embryos which produce secondary embryos are then excised and placed into a suitable liquid medium. After repeated selection for clusters of somatic embryos which multiplied as early, globular staged embryos, the suspensions are maintained as described below.

Soybean embryogenic suspension cultures can be maintained in 35 mL liquid media on a rotary shaker, 150 rpm, at 26°C with fluorescent lights on a 16:8 hour day/night schedule. Cultures are subcultured every two weeks by inoculating approximately 35 mg of tissue into 35 mL of liquid medium.

Soybean embryogenic suspension cultures may then be transformed by the method of particle gun bombardment (Klein et al. (1987) *Nature* (London) 327:70-73, U.S. Patent No. 4,945,050). A DuPont Biolistic™ PDS1000/HE instrument (helium retrofit) can be used for these transformations.

5 A selectable marker gene which can be used to facilitate soybean transformation is a chimeric gene composed of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812), the hygromycin phosphotransferase gene from plasmid pJR225 (from *E. coli*; Gritz et al. (1983) *Gene* 25:179-188) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*. The seed expression
10 cassette comprising the phaseolin 5' region, the fragment encoding the instant polypeptides and the phaseolin 3' region can be isolated as a restriction fragment. This fragment can then be inserted into a unique restriction site of the vector carrying the marker gene.

To 50 µL of a 60 mg/mL 1 µm gold particle suspension is added (in order): 5 µL DNA (1 µg/µL), 20 µl spermidine (0.1 M), and 50 µL CaCl₂ (2.5 M). The particle
15 preparation is then agitated for three minutes, spun in a microfuge for 10 seconds and the supernatant removed. The DNA-coated particles are then washed once in 400 µL 70% ethanol and resuspended in 40 µL of anhydrous ethanol. The DNA/particle suspension can be sonicated three times for one second each. Five µL of the DNA-coated gold particles are then loaded on each macro carrier disk.

20 Approximately 300-400 mg of a two-week-old suspension culture is placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5-10 plates of tissue are normally bombarded. Membrane rupture pressure is set at 1100 psi and the chamber is evacuated to a vacuum of 28 inches mercury. The tissue is placed approximately 3.5 inches away from the
25 retaining screen and bombarded three times. Following bombardment, the tissue can be divided in half and placed back into liquid and cultured as described above.

Five to seven days post bombardment, the liquid media may be exchanged with fresh media, and eleven to twelve days post bombardment with fresh media containing 50 mg/mL hygromycin. This selective media can be refreshed weekly. Seven to eight weeks post
30 bombardment, green, transformed tissue may be observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue is removed and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Each new line may be treated as an independent transformation event. These suspensions can then be subcultured and maintained as clusters of immature embryos or
35 regenerated into whole plants by maturation and germination of individual somatic embryos.

EXAMPLE 7

Expression of Chimeric Genes in Microbial Cells

The cDNAs encoding the instant polypeptides can be inserted into the T7 *E. coli* expression vector pBT430. This vector is a derivative of pET-3a (Rosenberg et al. (1987) *Gene* 56:125-135) which employs the bacteriophage T7 RNA polymerase/T7 promoter system. Plasmid pBT430 was constructed by first destroying the EcoR I and Hind III sites in pET-3a at their original positions. An oligonucleotide adaptor containing EcoR I and Hind III sites was inserted at the BamH I site of pET-3a. This created pET-3aM with additional unique cloning sites for insertion of genes into the expression vector. Then, the Nde I site at the position of translation initiation was converted to an Nco I site using oligonucleotide-directed mutagenesis. The DNA sequence of pET-3aM in this region, 5'-CATATGG, was converted to 5'-CCCATGG in pBT430.

Plasmid DNA containing a cDNA may be appropriately digested to release a nucleic acid fragment encoding the protein. This fragment may then be purified on a 1% NuSieve GTG™ low melting agarose gel (FMC). Buffer and agarose contain 10 µg/ml ethidium bromide for visualization of the DNA fragment. The fragment can then be purified from the agarose gel by digestion with GELase™ (Epicentre Technologies) according to the manufacturer's instructions, ethanol precipitated, dried and resuspended in 20 µL of water. Appropriate oligonucleotide adapters may be ligated to the fragment using T4 DNA ligase (New England Biolabs, Beverly, MA). The fragment containing the ligated adapters can be purified from the excess adapters using low melting agarose as described above. The vector pBT430 is digested, dephosphorylated with alkaline phosphatase (NEB) and deproteinized with phenol/chloroform as described above. The prepared vector pBT430 and fragment can then be ligated at 16°C for 15 hours followed by transformation into DH5 electrocompetent cells (GIBCO BRL). Transformants can be selected on agar plates containing LB media and 100 µg/mL ampicillin. Transformants containing the gene encoding the instant polypeptides are then screened for the correct orientation with respect to the T7 promoter by restriction enzyme analysis.

For high level expression, a plasmid clone with the cDNA insert in the correct orientation relative to the T7 promoter can be transformed into *E. coli* strain BL21(DE3) (Studier et al. (1986) *J. Mol. Biol.* 189:113-130). Cultures are grown in LB medium containing ampicillin (100 mg/L) at 25°C. At an optical density at 600 nm of approximately 1, IPTG (isopropylthio-β-galactoside, the inducer) can be added to a final concentration of 0.4 mM and incubation can be continued for 3 h at 25°. Cells are then harvested by centrifugation and re-suspended in 50 µL of 50 mM Tris-HCl at pH 8.0 containing 0.1 mM DTT and 0.2 mM phenyl methylsulfonyl fluoride. A small amount of 1 mm glass beads can be added and the mixture sonicated 3 times for about 5 seconds each time with a microprobe sonicator. The mixture is centrifuged and the protein concentration of the supernatant

EXAMPLE 8

**Evaluating Compounds for Their Ability to Inhibit the Activity
of Vitamin E Biosynthetic Enzymes**

Purification of the instant polypeptides, if desired, may utilize any number of separation technologies familiar to those skilled in the art of protein purification. Examples of such methods include, but are not limited to, homogenization, filtration, centrifugation, heat denaturation, ammonium sulfate precipitation, desalting, pH precipitation, ion exchange chromatography, hydrophobic interaction chromatography and affinity chromatography, wherein the affinity ligand represents a substrate, substrate analog or inhibitor. When the instant polypeptides are expressed as fusion proteins, the purification protocol may include the use of an affinity resin which is specific for the fusion protein tag attached to the expressed enzyme or an affinity resin containing ligands which are specific for the enzyme. For example, the instant polypeptides may be expressed as a fusion protein coupled to the C-terminus of thioredoxin. In addition, a (His)₆ peptide may be engineered into the N-terminus of the fused thioredoxin moiety to afford additional opportunities for affinity purification. Other suitable affinity resins could be synthesized by linking the appropriate ligands to any suitable resin such as Sepharose-4B. In an alternate embodiment, a thioredoxin fusion protein may be eluted using dithiothreitol; however, elution may be accomplished using other reagents which interact to displace the thioredoxin from the resin. These reagents include β -mercaptoethanol or other reduced thiol. The eluted fusion protein may be subjected to further purification by traditional means as stated above, if desired. Proteolytic cleavage of the thioredoxin fusion protein and the enzyme may be accomplished

after the fusion protein is purified or while the protein is still bound to the ThioBond™ affinity resin or other resin.

Crude, partially purified or purified enzyme, either alone or as a fusion protein, may be utilized in assays for the evaluation of compounds for their ability to inhibit enzymatic activation of the instant polypeptides disclosed herein. Assays may be conducted under well known experimental conditions which permit optimal enzymatic activity. For example, assays for gamma tocopherol methyltransferase are presented by d'Harlingue and Camara (1985) *J. Biol. Chem.* 260:15200-15203. Assays for 4-hydroxyphenylpyruvate dioxygenase are presented by Norris et al. (1995) *Plant Cell* 7: 2139-2149.

EXAMPLE 9

Expression of gamma Tocopherol Methyltransferase in Soybean Somatic Embryos

The ability to change the levels of alpha- and gamma-tocopherol in plants by transforming them with sequences encoding gamma tocopherol methyltransferase was tested by preparing transgenic soybean somatic embryos and assaying the tocopherol levels.

Plasmid DNA from clone sah1c.pk001.k8 was used as a template to prepare a Not I pcr fragment encoding the entire deduced open reading frame using the following pcr primers (forward primer AGC GCG GCC GCA TGG CCA CCG TGG TGA GGA TCC CAA CAA TCT CAT GCA TCC ACA TCC ACA; reverse primer AGC GCG GCC GCT TAT CTA

GTG TGG AAA TAA TGA TCA). Standard pcr reactions were used (100 microliter total reaction containing 5 ng plasmid, 25 nmoles primer, 25 nmoles dNTPs, 1x cloned pfu buffer (Stratagene), 5% DMSO, 5U cloned pfu DNA polymerase (Stratagene). The recommended cycling parameters for pcr with pfu DNA polymerase were used (denature 45 s 94°C, anneal 45 s 55°C, extend 2.5 min 72°C, 25 cycles, final 10min extension at 72°C). The pcr product was purified on a 1% agarose/TAE gel (precast FMC), the ethidium bromide visualized band cut out and purified using a QIAquick gel extraction kit (Qiagen). The band prep (40 ng) was ligated into PCR Blunt (InVitrogen) according to the manufacturers recommendations, and the ligated plasmid used to transform *E. coli* DH10 cells. Kanamycin resistant colonies were grown overnight in liquid culture (LB/Kan), plasmids prepared and cut with Not I.

Plasmids containing the correct insert size were selected for full insert sequence to confirm fidelity of pcr. Plasmids containing the correct insert verified by DNA sequencing were digested with Not I and ligated to Not I-digested and phosphatase-treated pKS67. The plasmid pKS67 was prepared by replacing in pRB20 (described in U.S. Patent No. 5,846,784) the 800 bp Nos 3' fragment, with the 285 bp Nos 3' fragment containing the polyadenylation signal sequence and described in Depicker et al. (1982) *J. Mol. Appl. Genet.* 1:561-573. Clones were screened for the sense and anti-sense orientation of the gamma tocopherol methyltransferase insert fragment by restriction enzyme digestion.

Transformation of Soybean Somatic Embryo Cultures

The following stock solutions and media were used for transformation and propagation of soybean somatic embryos:

Stock Solutions		Media
<u>MS Sulfate 100x stock</u> (g/L)		<u>SB55 (per Liter)</u>
MgSO ₄ ·7H ₂ O	37.0	10 mL of each MS stock
MnSO ₄ ·H ₂ O	1.69	1 mL of B5 Vitamin stock
ZnSO ₄ ·7H ₂ O	0.86	0.8 g NH ₄ NO ₃
CuSO ₄ ·5H ₂ O	0.0025	3.033 g KNO ₃
		1 mL 2,4-D (10 mg/mL stock)
		0.667 g asparagine
		pH 5.7
<u>MS Halides 100x stock</u>		<u>SB103 (per Liter)</u>
CaCl ₂ ·2H ₂ O	44.0	1 pk. Murashige & Skoog salt mixture*
KI	0.083	60 g maltose
CoCl ₂ ·6H ₂ O	0.00125	2 g gelrite
KH ₂ PO ₄	17.0	pH 5.7
H ₃ BO ₃	0.62	
Na ₂ MoO ₄ ·2H ₂ O	0.025	
Na ₂ EDTA	3.724	
FeSO ₄ ·7H ₂ O	2.784	
		<u>SB148 (per Liter)</u>
<u>B5 Vitamin stock</u>		1 pk. Murashige & Skoog salt mixture*
myo-inositol	100.0	60 g maltose
nicotinic acid	1.0	1 mL B5 vitamin stock
pyridoxine HCl	1.0	7 g agarose
thiamine	10.0	pH 5.7

5

*(Gibco BRL)

Soybean embryonic suspension cultures were maintained in 35 mL liquid media (SB55) on a rotary shaker (150 rpm) at 28°C with a mix of fluorescent and incandescent lights providing a 16 h day 8 h night cycle. Cultures were subcultured every 2 to 3 weeks by inoculating approximately 35 mg of tissue into 35 mL of fresh liquid media.

Soybean embryonic suspension cultures were transformed with the plasmid containing the gamma tocopherol methyltransferase sequence by the method of particle gun bombardment (see Klein et al. (1987) *Nature* 327:70-73) using a DuPont Biolistic PDS1000/He instrument. Five µL of pKS93s plasmid DNA (1 g/L), 50 µL CaCl₂ (2.5 M), and 20 µL spermidine (0.1 M) were added to 50 µL of a 60 mg/mL 1 mm gold particle suspension. The particle preparation was agitated for 3 minutes, spun on a microfuge for

10 seconds and the supernate removed. The DNA-coated particles were then washed once with 400 μ L of 70% ethanol and resuspended in 40 μ L of anhydrous ethanol. The DNA/particle suspension was sonicated three times for 1 second each. Five μ L of the DNA-coated gold particles were then loaded on each macro carrier disk.

- 5 Approximately 300 to 400 mg of two-week-old suspension culture was placed in an empty 60 mm X 15 mm petri dish and the residual liquid removed from the tissue using a pipette. The tissue was placed about 3.5 inches away from the retaining screen and bombarded twice. Membrane rupture pressure was set at 1100 psi and the chamber was evacuated to -28 inches of Hg. Two plates were bombarded, and following bombardment, 10 the tissue was divided in half, placed back into liquid media, and cultured as described above.

- 15 Fifteen days after bombardment, the liquid media was exchanged with fresh SB55 containing 50 mg/mL hygromycin. The selective media was refreshed weekly. Six weeks after bombardment, green, transformed tissue was isolated and inoculated into flasks to generate new transformed embryonic suspension cultures.

- 20 Transformed embryonic clusters were removed from liquid culture media and placed on a solid agar media, SB103, containing 0.5% charcoal to begin maturation. After 1 week, embryos were transferred to SB103 media minus charcoal. After 5 weeks on SB103 media, maturing embryos were separated and placed onto SB148 media. During maturation 20 embryos were kept at 26°C with a mix of fluorescent and incandescent lights providing a 16 h day 8 h night cycle. After 3 weeks on SB148 media, embryos were analyzed for the expression of the tocopherols. Each embryonic cluster gave rise to 5 to 20 somatic embryos.

Non-transformed somatic embryos were cultured by the same method as used for the transformed somatic embryos.

- 25 *Analysis of Transformed Somatic Embryos*

- 30 At the end of the 6th week on SB148 medium somatic embryos were harvested from 25 independently transformed lines. Somatic embryos were collected in pools of five and weighed for fresh weight. Excess embryos were stored in 96-well plates at -80°. The pooled somatic embryos were lyophilized for 18 hours and the dry weight measured. The lyophilized somatic embryos were briefly pulverised with a hand held Potter homogeniser and then 600 μ L of heptane added and the samples incubated for 24 hours in the dark at room temperature to extract oils and tocopherols. The heptane was decanted and a further 300 μ L added to the samples. The extracts were combined and centrifuged (5 min, 12000 g). The supernatant was stored in amber hplc autosampler vials at -20°C prior to analysis.

- 35 HPLC analysis of the extracts was carried out using an HP1100 system (Agilent Technologies) 25 μ L of the heptane sample was applied to a Lichrosphere Si 60 column (5 micron 4 x 12.5 mm). The column was eluted with heptane/isopropanol (98:2 v/v) at a flow rate of 1 mL/min. After 6 minutes all four tocopherol isomers were eluted, as detected by

a HP1100 fluorescence detector (Excitation wavelength 295 nm, emission wavelength 330 nm). Individual tocopherol standards (Matreya) were diluted with hplc grade heptane to levels between 1 and 200 ng/ul to construct a 6 point external standard curve. Tocopherols in each oil were quantified using a standard curve run on the same day as the samples. The

5 sum of tocopherol peak areas of samples from a non-transformed control line were compared with those of 25 independent gamma tocopherol methyltransferase-transformed, hygromycin resistant lines. Figure 3 shows a graph depicting the distribution of the percent alpha- and gamma-tocopherol of soybean somatic embryos transgenic for the gamma tocopherol methyltransferase gene and a control line. The mean alpha tocopherol content is 40 to 70%.

10 Some lines, such as the ones represented in bars 2, 5, and 18 appear to represent overexpression of the gamma tocopherol methyltransferase, which would be predicted to yield higher levels of alpha tocopherol. Other lines, such as those represented in bars 6 and 10 appear to represent co-suppression of the gamma tocopherol methyltransferase since this is predicted to yield higher levels of gamma tocopherol. These results indicate that

15 transgenic expression of gamma-tocopherol methyltransferase affords the ability to manipulate tocopherol levels as desired for a particular application.

Various modifications of the invention in addition to those shown and described herein will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

20 The disclosure of each reference set forth above is incorporated herein by reference in its entirety.

CLAIMS

What is claimed is:

1. An isolated polynucleotide comprising a first nucleotide sequence encoding a polypeptide of at least 160 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8, 10, 22, 24, 26, 28, and 30, or a second nucleotide sequence comprising the complement of the first nucleotide sequence.
2. An isolated polynucleotide comprising a first nucleotide sequence encoding a polypeptide of at least 150 amino acids that has at least 95% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:12, 14, 16, 18, 20, 32, 34, 36, and 38, or a second nucleotide sequence comprising the complement of the first nucleotide sequence.
3. The isolated polynucleotide of Claim 1 or Claim 2, wherein the first nucleotide sequence consists of a nucleic acid sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, and 37 that codes for the polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, and 38.
4. The isolated polynucleotide of Claim 1 or Claim 2 wherein the nucleotide sequences are DNA.
5. The isolated polynucleotide of Claim 1 or Claim 2 wherein the nucleotide sequences are RNA.
6. A chimeric gene comprising the isolated polynucleotide of Claim 1 or Claim 2 operably linked to suitable regulatory sequences.
7. An isolated host cell comprising the chimeric gene of Claim 6.
8. A host cell comprising an isolated polynucleotide of Claim 1 or Claim 2.
9. The host cell of Claim 8 wherein the host cell is selected from the group consisting of yeast, bacteria, plant, and virus.
10. A virus comprising the isolated polynucleotide of Claim 1 or Claim 2.
11. A polypeptide of at least 160 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8, 10, 22, 24, 26, 28, and 30.
12. A polypeptide of at least 150 amino acids that has at least 95% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:12, 14, 16, 18, 20, 32, 34, 36, and 38.
13. A method of selecting an isolated polynucleotide that affects the level of expression of a vitamin E biosynthetic enzyme polypeptide in a plant cell, the method comprising the steps of:

(a) constructing an isolated polynucleotide comprising a nucleotide sequence of at least one of 30 contiguous nucleotides derived from an isolated polynucleotide of Claim 1 or Claim 2;

(b) introducing the isolated polynucleotide into a plant cell; and

5 (c) measuring the level of a polypeptide in the plant cell containing the polynucleotide to provide a positive selection means.

14. The method of Claim 13 wherein the isolated polynucleotide consists of a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, and 37 that codes for the polypeptide selected from
10 the group consisting of SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, and 38.

15 15. A method of selecting an isolated polynucleotide that affects the level of expression of a vitamin E biosynthetic enzyme polypeptide in a plant cell, the method comprising the steps of:

(a) constructing an isolated polynucleotide of Claim 1 or Claim 2;

(b) introducing the isolated polynucleotide into a plant cell; and

(c) measuring the level of polypeptide in the plant cell containing the polynucleotide to provide a positive selection means.

16. A method of obtaining a nucleic acid fragment encoding a vitamin E
20 biosynthetic enzyme polypeptide comprising the steps of:

(a) synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 30 contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, and the complement of such nucleotide sequences; and

25 (b) amplifying a nucleic acid sequence using the oligonucleotide primer.

17. A method of obtaining a nucleic acid fragment encoding a vitamin E biosynthetic enzyme polypeptide comprising the steps of:

(a) probing a cDNA or genomic library with an isolated polynucleotide comprising at least one of 30 contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37 and the complement of such nucleotide sequences;

(b) identifying a DNA clone that hybridizes with the isolated polynucleotide;

(c) isolating the identified DNA clone; and

35 (d) sequencing the cDNA or genomic fragment that comprises the isolated DNA clone.

18. A method for evaluating at least one compound for its ability to inhibit the activity of a vitamin E biosynthetic enzyme, the method comprising the steps of:

(a) transforming a host cell with a chimeric gene comprising a nucleic acid fragment encoding a vitamin E biosynthtetic enzyme, operably linked to suitable regulatory sequences;

5 (b) growing the transformed host cell under conditions that are suitable for expression of the chimeric gene wherein expression of the chimeric gene results in production of the vitamin E biosynthtetic enzyme encoded by the operably linked nucleic acid fragment in the transformed host cell;

(c) optionally purifying the vitamin E biosynthtetic enzyme expressed by the transformed host cell;

10 (d) treating the vitamin E biosynthtetic enzyme with a compound to be tested; and

(e) determining the activity of the vitamin E biosynthtetic enzyme that has been treated with the compound, thereby selecting compounds with potential for inhibitory activity.

15 19. A composition comprising the isolated polynucleotide of Claim 1 or Claim 2.

20. A composition comprising the isolated polypeptide of Claim 11.

21. An isolated polynucleotide comprising the nucleotide sequence having at least one of 30 contiguous nucleotides derived from a nucleic acid sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 20 35, 37 and the complement of such sequences.

22. An expression cassette comprising an isolated polynucleotide of Claim 1 or Claim 2 operably linked to a promoter.

23. A method for positive selection of a transformed cell comprising:

25 (a) transforming a host cell with an expression cassette of Claim 22; and (b) growing the transformed host cell under conditions which allow expression of the polynucleotide in an amount sufficient to complement a null mutant to provide a positive selection means.

24. The method of Claim 23 wherein the plant cell is a monocot.

25. The method of Claim 23 wherein the plant cell is a dicot.

30

FIGURE 1

```

SEQ ID NO:39  MVYHVRPKHAL-----FLAFYCYFS-----
SEQ ID NO:40  MKATLAAPSSLTSL-P-----YRTNSS-FGSKSLLFRSPSSSSVS-MTTTRGNV
SEQ ID NO:38  MAN-----SXRPA-PLTPLHRLDAAPPRLPSLGHAAARPVPRPVLPLPAR-XLRAPDGV
SEQ ID NO:28  MATVVRI-----PTISCIHIHTFRSQSPRTFARIUVGPRSWAPIRASA-ASSERGEI
SEQ ID NO:08  MATVVRI-----PTISCIHIHTFRSQSPRTFARIUVGPRSWAPIRASA-ASSERGEI
SEQ ID NO:22  MAHAALLHCSQSSR-SLAACRRGSHYRAPH-VPRHSRRLRRVVSL---R-PMASSTA-
SEQ ID NO:24  AR--VQPTGALAPLHPLLRCTSRHLCASASPRAGLCLHHHRRRRSSRRTKLAVRAMAPT
SEQ ID NO:26  FRH-----GH-----AL-----
1
60

SEQ ID NO:39  LLTMASATIASADLYEKIKNFYDDSSGLWEDVWGEHMHGYYGPHGTYRI---DRRQAI
SEQ ID NO:40  AVAAAATSTEALRKGIAEFYNETSGLWEEIWGDHMHGFDPSVQLSDSGHKEAQI
SEQ ID NO:38  VDDRGPGDAAPPGLKEGIAGLYDESSGLWESIWGEHMHGFDSEAAASMSD--HRRRAQI
SEQ ID NO:28  VLEQPKKDDKKLQKGIAEFYDESSGLWENIWGDHMHGFDSDTVSLSD--HRAAQI
SEQ ID NO:08  VLEQPKKDDKKLQKGIAEFYDESSGLWENIWGDHMHGFDSDTVSLSD--HRAAQI
SEQ ID NO:22  ---QAPATAPPGLKEGIAGLYDESSGLWENIWGDHMHGFDSEAAASMSD--HRRRAQI
SEQ ID NO:24  LSSSSTAAAAPPGLKEGIAGLYDESSGVWESIWGEHMHGFDAGEAASMSD--HRRRAQI
61
120

SEQ ID NO:39  DLIKELLAWAVPQNSA---KPRKILDGCGIGGSSLYLAQQHQAQAEVMGASLSPVQVERA
SEQ ID NO:40  RMIEESLRFAGVTDEEE-EKKIKKVVVDVCGGIGGSSRYLASKFGAECIGITLSPVQAKRA
SEQ ID NO:38  RMIEEALAFAAVP--DDPTNKPTIVDVCGGIGGSSRYLANKYGAQCSGITLSPVQAERG
SEQ ID NO:28  RMIQESLRFASV--SEERSKWPKSIVDVCGGIGGSSRYLAKKFGATSVGITLSPVQAQRA
SEQ ID NO:08  RMIQESLRFASV--SEERSKWPKSIVDVCGGIGGSSRYLAKKFGATSVGITLSPVQAQRA
SEQ ID NO:22  RMIEEALAFAGVPASDDPEKTPKTIVDVCGGIGGSSRYLAKKYGXQCTGITLSPVQAERG
SEQ ID NO:24  RMIEESLAF-----
121
180

```

FIGURE 1

```

SEQ ID NO:39 GERARALGLSTCQFQVANALDLPFASDSFQVWWSLESGEHMPNKAQFLQEAWRVLKPGG
SEQ ID NO:40 NDIAAQAQSLSHKASFQVADALDQPFEDGKFDLVWSMESGEHMPDKAKFVKELVRVAAPGG
SEQ ID NO:38 NALAAAQGLSDKASFQVADALEQPFDPGQFDLVWSMESGEHMPNKKQFVSELARVAAPGA
SEQ ID NO:28 NALAAAQGLADKVSFQVADALQPFSDGQFDLVWSMESGEHMPDKAKFVGELARVAAPGA
SEQ ID NO:08 NALAAAQGLADKVSFQVADALQPFSDGQFDLVWSMESGEHMPDKAKFVGELARVAAPGA
SEQ ID NO:22 NALAAAQGLSDQVTLQVADALEQPFDPGQFDLVWSMESGEHMPDKRKQFVSELARVAAPGG
SEQ ID NO:24 -----
SEQ ID NO:26 -----AQPFPDGQFDLVWSMESDEHMPDKRKQFVSELARVAAPGA
181 240

SEQ ID NO:39 RLILATWCHRPIDPGNGPLTADERRHLQAIYDVYCLPYVVSPLDYEAIARECGFGEIKTA
SEQ ID NO:40 RIIIVTWCHRNLSAGEEALQWEQNILDKICKTFYLPAWCSTDDYVNLQSHSLQDIKCA
SEQ ID NO:38 TIIIVTWCHRNLPASEDSLKPDENLLKKICDAYYLPDWCSPSDYVVKIAESLSLEDIKTA
SEQ ID NO:28 IIIIVTWCHRD LGPDEQSLHPWEQD LKKICDAYYLPAWCSTSDYVKKLLQSLSLQDIKSE
SEQ ID NO:08 IIIIVTWCHRD LGPDEQSLHPWEQD LKKICDAYYLPAWCSTSDYVKKLLQSLSLQDIKSE
SEQ ID NO:22 TIIIVTWCHRNLDPSETSLKPDELSLLRRICDAYYLPDWCSPSDYVVKIAESLSLEDIKTA
SEQ ID NO:24 -----
SEQ ID NO:26 RIIIVTWCHRNLEPSEESLKPDENLLKRICDAYYLPDWCSPSDYVVKIAESLSLEDIRTA
241 300

SEQ ID NO:39 DWSVAVAPFWDRVIESAFDPRVLWALGOAGPKIINAAALCLRLMKWGYERGLVRFGLLTGI
SEQ ID NO:40 DWSENVAPFWPAVIRALTATWGLVSLLRSGMKSIIKGALTMPLMIEGYKKGVIKFGIITCQ
SEQ ID NO:38 DWSENVAPFWPAVIRSAFTWGLTSLLRSGWKTIKGALVMPMLIQGYKKGLIKFSIITCR
SEQ ID NO:28 DWSRFVAPFWPAVIRSAFTWGLSLLSSGQKTIKGALAMPLMIEGYKKDLIKFAIITCR
SEQ ID NO:08 DWSRFVAPFWPAVIRSAFTWGLSLLSSGKLGII-----YIAFQKQTPPSSIATCK
SEQ ID NO:22 DWSENVAPFWPAVIRSAFTWGLTSLTGWKTIRGAMVMPLMIQGYKKGLIKFTIITCR
SEQ ID NO:24 -----
SEQ ID NO:26 DWSENVAPFWPAVIRSAFTWGLTSLLRSGWETVRGAMVMPLVIEGYKKGLIKFPIITCR
301 360

```

FIGURE 1

```
SEQ ID NO:39 KPLV-----  
SEQ ID NO:40 KPL-----  
SEQ ID NO:38 KPQAAIEGEPEEAAPSVE.-  
SEQ ID NO:28 K-----PE.-  
SEQ ID NO:08 SYVTDHYFHTR.-  
SEQ ID NO:22 KPGAA-.MA---HAALLHC  
SEQ ID NO:24 -----  
SEQ ID NO:26 KPETT-----Q.-  
361 380
```

FIGURE 2

SEQ ID NO: 41	MPPTPTTAAATGAAAAVTPEHARPH-----RMVRFNPRSDREHTLSFHHVEFWCADAASA	60
SEQ ID NO: 43	MGHQNAAVSENQNHDDGAASSPGFKLVGFSGKFKVRKNPKSDKEKVKRFHHIEFWCGDATNV	
SEQ ID NO: 42	MGKKQSEAEILSS-NSSNTSPATFKLVGFNNFVRANPKSDHEFAVKRFHHIEFWCGDATNT	
SEQ ID NO: 36	MPIPMCNEIQQA-QAQAQAQPGFKLVGFKNFVRTNPKSDRFQVNRFFHHIEFWCTDATNA	
SEQ ID NO: 38	MPPTPTTAAATGAAA-VTPEHARPR-----RMVRFNPRSDREHTLAFHHVEFWCADAASA	
SEQ ID NO: 32	MGK-QTTTSATAA-DGSKDAHAEFKLVGFKNFVRTNPKSDHFCVHREFFHHIEFWCGDATNT	1
		60
SEQ ID NO: 41	AGRFALGAPLAARSDLSTGNSAHASQLLRSGSLAFLFTAPYANG-----CDAATASL	
SEQ ID NO: 43	ARRFSWGLGMRFSAKSDLSTGNMVMHASYLSTSGDLRFLFTAPYSPSLSAGEIKPTTTASI	
SEQ ID NO: 42	SRRFSWGLGMLVAKSDLSTGNSVHASYLVRSANLSFVFTAPYSPSTTSS-G----SAAI	
SEQ ID NO: 36	SRRFSWGLGMPIVAKSDLSTGNQIHASYLRLSGDLSFLFSAPYSPSLSAGS-SAASSASI	
SEQ ID NO: 38	AGRFALGAPLAARSDLSTGNSVHASQLLRSGNLAFLFTAPYANG-----CDAATASL	
SEQ ID NO: 32	AKRFSWGLGMLVAKSDLSTGNSAHASYLRLSGELNFLTSPYSPSISAPS-----SAAI	61
		120
SEQ ID NO: 41	PSFSADAAARRFSADHGIAVRSVALRVADAAEAFRRRRRGARPAFAPVDLGRGFA-FAEV	
SEQ ID NO: 43	PSFDHGSCRSFFSSHGLGVRAVAIEVEDAESAFSISVANGAIPSSPPIVLNEAVT-IAEV	
SEQ ID NO: 42	PSFSASGFHSFAAKHGLAVRAIAIEVADVAAAFASVARGARPASAPVEL-DDQAWLAEV	
SEQ ID NO: 36	PSFDAATCLAFAAKHGFGVRAIAIEVADAEAAFSASVAKGAEPASPPV-LVDDRTGFAEV	
SEQ ID NO: 38	PSFSADAAARQFSADHGLAVRSIALRVADAAEAFRASVDGGARPAFSPVDLGRGFG-FAEV	
SEQ ID NO: 32	PSFSFSTYQSFSTSSHGLAVRAVAIQVDSAFSAYSASISRGAKPVSAPILLSDNKTIAIEV	121
		180

FIGURE 2

```

SEQ ID NO:41  ELYGDVVLRFVSH-PDGTDPV-----FLPGFEGVTNPDA---VDYGLTRFDHVVGNVNVP      240
SEQ ID NO:43  KLYGDVVLRFVSYKAEDEKSE-----FLPGFERVEDASSFP-LDYGIRRLDHAVGNVP
SEQ ID NO:42  ELYGDVVLRFVSFGRE-----EGLFLPGFEAVEGTASFDPDLDYGIRRLDHAVGNVT
SEQ ID NO:36  RLYGDVVLRFVSYKDAAPQAPHADPSRWFLPGFEAAAASSSSFPELDYGIRRLDHAVGNVP
SEQ ID NO:38  ELYGDVVLRFVSH-PDGRDVP-----FLPGFEGVSNPDA---VDYGLTRFDHVVGNVNVP
SEQ ID NO:32  HLYGDSVLRVFSYGDNG-----TGPDGWFLPGFEPVDDQMSYKELDYGIRRLDHAVGNVP      240
181

SEQ ID NO:41  ELAPAAAYIAGFTGFHEFAEFTAEADVGTTESGLNSVVLANNSEGVLLPLNEPVHGTKRRS
SEQ ID NO:43  ELGPALTYVAGFTGFHQFAEFTADDVGTAEGLNSAVLASNDEMVLPLINEPVHGTKRKS
SEQ ID NO:42  ELGPVVEYIKFTGFHEFAEFTAEADVGTLEGLNSVVLANNSEEMVLLPLNEPVYGTKRKS
SEQ ID NO:36  ELAPAVRYLKGFSGFHEFAEFTAEADVGTSEGLNSVVLANNSETVLLPLNEPVYGTKRKS
SEQ ID NO:38  ELAPAAAYVAGFTGFHEFAEFTEDVGTAEGLNSMVLANNSEGVLLPLNEPVHGTKRRS
SEQ ID NO:32  ELGPVVDYLLKKFTGFHEFAEFTSEDVGTAEGLNSMVLANNSEGVLLPLNEPVFGTKRKS      300
241

SEQ ID NO:41  QIQTFLEHHGGPGVQHIAVASSDVLRTLRLKMRARSAMGGFDLPPPLPKYYEGVRRLAGD
SEQ ID NO:43  QIQTYLEHNEGAGLQHLALMSEDI FRTLREMRKRSSIGGDFDMPSPPPTYYQNLKRVGD
SEQ ID NO:42  QIQTYLEHNEGAGVQHLALVSEDI FRTLREMRKRSClGGFEFMPSPPTYYKNLKNRVGD
SEQ ID NO:36  QIETYLEHNEGAGVQHLALVTHDI FRTLREMRKRSLGGFEFMPSPPTYYANLHNRAAD
SEQ ID NO:38  QIQTFLEHHGGSGVQHIAVASSDVLRTLREMRARSAMGGFDLPPPLPKYYEGVRRRIAGD
SEQ ID NO:32  QIQTYLEHNEGPGVQHLALVSEDI FNTLREMRKRSGVGGFEFMPSPPLTYYKNLKNRAGD      360
301

```


FIGURE 2

```

SEQ ID NO:41  VLSEAQIKECQELGVLVDRDDQGVLLQIFTKPVGDRPTLFLEMIQIRIGCMKDERGEEYQ
SEQ ID NO:43  VLSDDQIKECEELGILVDRDDQGTLLQIFTKPLGDRPTIFIEIIQIRVGCMMKDEEGKAYQ
SEQ ID NO:42  VLSDEQIKECEDLGILVDRDDQGTLLQIFTKPVGDRPTLFIEIIQIRVGCMLKDDAGQMYQ
SEQ ID NO:36  VLTVDQIKQCEELGILVDRDDQGTLLQIFTKPVGDRPTIFIEIIQIRIGCMVEDEEGKVYQ
SEQ ID NO:38  VLSEAQIKECQELGVLVDRDDQGVLLQIFTKPVGDRPTLFLEMIQIRIGCMKDERGEEYQ
SEQ ID NO:32  VLRDEQIEECEKLGILVDRDDQGTLLQIFTKPVGDRPTLFIEIIQIRIGCMKDEQGKLYQ
361
SEQ ID NO:41  KGGCGGFGKGNFSELFKSIEDYEKSLEAKQSAAVQG-S
SEQ ID NO:43  SGGCGGFGKGNFSELFKSIEEYEKTLQAKQLVG-----
SEQ ID NO:42  KGGCGGFGKGNFSELFKSIEEYEKTLQAKQITGSA-AA
SEQ ID NO:36  KGACGGFGKGNFSELFKSIEEYEKTLQAKR-----TA.
SEQ ID NO:38  KGGCGGFGKGNFSELFKSIEDYEKSLEAKQSAAVQGS.
SEQ ID NO:32  KSGCGGFGKGNFSELFKSIEEYEKMLEAKQVTETASA.
421
420
458

```


Figure 3



SEQUENCE LISTING

<110> Rafalski, J. Antoni
 Cahoon, Rebecca E.
 Coughlan, Sean
 Miao, Guo-Hua

<120> PLANT VITAMIN E BIOSYNTHETIC ENZYMES

<130> BB1289

<140>

<141>

<150> 60/110,781

<151> 1998-12-03

<160> 43

<170> Microsoft Office 97

<210> 1

<211> 792

<212> DNA

<213> Zea mays

<400> 1

```

gttgagccct gttcaagccg agagaggaaa tgctctcgct gcagcgcagg ggttgtcgga 60
tcagggttact ctgcaagttg ctgatgctct ggagcaaccg ttccctgacg ggcagttcga 120
tctgggtgtgg tccatggaga gtggcgagca catgccggac aagagaaaagt ttgttagtga 180
gctagcacgc gtggcggtc ctggaggagc aataatcatc gtgacatggt gccataggaa 240
cctggatcca tccgaaacct cgctaaagcc cgatgaactg agcctcctga ggaggatatg 300
cgacgcgtac tacctcccgg actggtgctc accttcagac tatgtgaaca ttgccaagtc 360
actgtctctc gaggatatca agacagctga ctggtcggag aacgtggccc cgttttggcc 420
cgccgtgata aaatcagcgc taacatggaa gggcttcacc tctctgctga cgaccggatg 480
gaagacgata agaggcgaga tggatgatgc gctaataatc cagggtctaca agaaggggct 540
catcaaattc accatcatca cctgtcgcaa gcctggagcc gcgtangagg aggcaaggag 600
cacaagttac tagcacagca caggatgcaa gtgcatatgt agatcatggc acatcgccgt 660
cactcatcat actgcacaaa atcaaattc caggacattt aataattctg cacctcanat 720
attcaggggg gccggtacca atcgccatat gatctatacc gnccacggcg tcnttaacte 780
tnacggaaan ct 792

```

<210> 2

<211> 191

<212> PRT

<213> Zea mays

<400> 2

```

Leu Ser Pro Val Gln Ala Glu Arg Gly Asn Ala Leu Ala Ala Ala Gln
1           5           10           15

```

```

Gly Leu Ser Asp Gln Val Thr Leu Gln Val Ala Asp Ala Leu Glu Gln
20           25           30

```

```

Pro Phe Pro Asp Gly Gln Phe Asp Leu Val Trp Ser Met Glu Ser Gly
35           40           45

```

```

Glu His Met Pro Asp Lys Arg Lys Phe Val Ser Glu Leu Ala Arg Val
50           55           60

```

Ala Ala Pro Gly Gly Thr Ile Ile Ile Val Thr Trp Cys His Arg Asn
65 70 75 80

Leu Asp Pro Ser Glu Thr Ser Leu Lys Pro Asp Glu Leu Ser Leu Leu
85 90 95

Arg Arg Ile Cys Asp Ala Tyr Tyr Leu Pro Asp Trp Cys Ser Pro Ser
100 105 110

Asp Tyr Val Asn Ile Ala Lys Ser Leu Ser Leu Glu Asp Ile Lys Thr
115 120 125

Ala Asp Trp Ser Glu Asn Val Ala Pro Phe Trp Pro Ala Val Ile Lys
130 135 140

Ser Ala Leu Thr Trp Lys Gly Phe Thr Ser Leu Leu Thr Thr Gly Trp
145 150 155 160

Lys Thr Ile Arg Gly Ala Met Val Met Pro Leu Met Ile Gln Gly Tyr
165 170 175

Lys Lys Gly Leu Ile Lys Phe Thr Ile Ile Thr Cys Arg Lys Pro
180 185 190

<210> 3
<211> 521
<212> DNA
<213> Oryza sativa

<400> 3
cttacagaca aacggcagtt tgtaagcgag ctggcacgcg tcgcagctcc tggggcgaga 60
ataatcattg tgacctggtg ccataggaac ctgcagccat ccgaagagtc cctgaaacct 120
gatgagctga atctcctgaa aaggatatgc gatgcatatt atctcccaga ctggtgctct 180
ccttctgatt atgtcaaaat tgccgagtca ctgtctcttg aggatataag gacagctgat 240
tggtcaagag aacgtcgccc caatccggnc tgcnggttat taaatnaagc aattgacatg 300
gnaagggtta actttctcct ggctaagaan tgggtgggaa gacgattaag aaggtggaat 360
gggtgatgcc tccggatgat nnaaggntac aaagaaangg gtcaacaaat ttaacaanaa 420
caacctgtnc caaagncccg aaacaacgca ataatacccc antaatnaaa ttncgctcct 480
ggctaaccct ctccaacaac gaattaatgg aaanttctga c 521

<210> 4
<211> 82
<212> PRT
<213> Oryza sativa

<400> 4
Leu Thr Asp Lys Arg Gln Phe Val Ser Glu Leu Ala Arg Val Ala Ala
1 5 10 15

Pro Gly Ala Arg Ile Ile Ile Val Thr Trp Cys His Arg Asn Leu Glu
20 25 30

Pro Ser Glu Glu Ser Leu Lys Pro Asp Glu Leu Asn Leu Leu Lys Arg
35 40 45

Ile Cys Asp Ala Tyr Tyr Leu Pro Asp Trp Cys Ser Pro Ser Asp Tyr
50 55 60

Val Lys Ile Ala Glu Ser Leu Ser Leu Glu Asp Ile Arg Thr Ala Asp
65 70 75 80

Trp Ser

<210> 5
 <211> 592
 <212> DNA
 <213> Oryza sativa

<400> 5
 cttacatgta agctcgtgcc gaattcggca cgagcttaca aaatggccca cgccgccgcg 60
 gccacgggag cactggcacc gctgcatcca ctgctccgct gcacgagccg tcatctctgc 120
 gcctcggctt cccctcgcgc cggcctctgc ctccaccacc accgccgccg ccgccgcagc 180
 agccggagga cgaaactcgc cgtgcgcgcg atggcaccga cgttgctctc gtcgtcgacg 240
 gggcgggcag ctccccggg gctgaaggag ggcacgcgcg ggctctacga cgaancgtcc 300
 ggcgtgtggg agagcatctg gggcgagcac atgcaccacg gnttctacga cgccggcgag 360
 ggccgctcca tgtccgacca ccgccgcgcc ccagttcgca tgatcgagga cctcgccttc 420
 gccgcctccc cgatgatcgg agaagaacca aaatgtattg atttggtgtg gattggtggt 480
 actcaagata ntgngaaca atacggacgc atgctacgna tacttgatcg gtgcaggatga 540
 aagagaaaanc ctgcgcgaga caaggtatag caagtcctt taagttgtat ga 592

<210> 6
 <211> 144
 <212> PRT
 <213> Oryza sativa

<400> 6
 Leu Thr Cys Lys Leu Val Pro Asn Ser Ala Arg Ala Tyr Lys Met Ala
 1 5 10 15
 His Ala Ala Ala Ala Thr Gly Ala Leu Ala Pro Leu His Pro Leu Leu
 20 25 30
 Arg Cys Thr Ser Arg His Leu Cys Ala Ser Ala Ser Pro Arg Ala Gly
 35 40 45
 Leu Cys Leu His His His Arg Arg Arg Arg Arg Ser Ser Arg Arg Thr
 50 55 60
 Lys Leu Ala Val Arg Ala Met Ala Pro Thr Leu Ser Ser Ser Ser Thr
 65 70 75 80
 Ala Ala Ala Ala Pro Pro Gly Leu Lys Glu Gly Ile Ala Gly Leu Tyr
 85 90 95
 Asp Glu Xaa Ser Gly Val Trp Glu Ser Ile Trp Gly Glu His Met His
 100 105 110
 His Gly Phe Tyr Asp Ala Gly Glu Gly Ala Ser Met Ser Asp His Arg
 115 120 125
 Arg Ala Pro Val Arg Met Ile Glu Asp Leu Ala Phe Ala Ala Ser Pro
 130 135 140

<210> 7
 <211> 1331
 <212> DNA
 <213> Glycine max

<400> 7

```

gtgacatggc caccgtggtg aggatcccaa caatctcatg catccacatc cacacgttcc 60
gttcccaatc cctcgcact ttcgccagaa tccgggtcgg acccaggtcg tgggtccta 120
ttcgggcacg ggcagcgagc tcggagagag gggagatagt attggagcag aagccgaaga 180
aggatgacaa gaagaagctg cagaagggaa tcgcagagtt ttacgacgag tcgtctggct 240
tatgggagaa catttggggc gaccacatgc accatggctt ttatgactcg gattccactg 300
tttcgctttc ggatcatcgt gctgctcaga tccgaatgat ccaagagtct cttcgctttg 360
cctctgtttc tgaggagcgt agtaaatggc ccaagagtat agttgatggt gggtgtggca 420
taggtggcag ctctagatac ctggccaaga aatttggagc aaccagtgtg ggcatactc 480
tgagtcctgt tcaagctcaa agagcaaagt ctcttgctgc tgctcaagga ttggctgata 540
agggtttcct tcaggttgct gacgctctac agcaaccatt ctctgacggc cagtttgatc 600
tggtgtggtc catggagagt ggagagcata tgccctgacaa agctaagttt gttggagagt 660
tagctcgggt agcagcacca ggtgccatta taataatagt aacatggtgc cacagggact 720
ttggccctga cgaacaatcc ttacatccat gggagcaaga tctcttaaag aagatttgcg 780
atgcatatta cctccctgcc tgggtctcaa cttctgatta tgtaagtgtg ctccaatccc 840
tgtcacttca ggacatcaag tcagaagatt ggtctcgctt tgttgctcca tttggccag 900
cagtgatacg ctcagccttc acatggaagg gtctatcttc actcttgagc agtggtgaagc 960
ttggaattta tattgcattt caaaaacaaa cccccccatc ttctattgca acttgcaagt 1020
cttatgtcac tgatcattat ttccacacta gataaccctt tacaactaag aacgtagtct 1080
tcattgttcag cgaaatagat aaaaatatgc aacagagtca gagacagggt gcatgatatt 1140
tacaagaaaa tatcttttat atatataaat gattcaatca aattacttga tgaggattat 1200
gagtgaanaa gagaggacag tcatagaaac tttatcctac attccttcta tttccacttc 1260
tgtcaaatat tcctttcatc ttagctatgc tacttgactt gagtaaaaaa aaaaaaaaaa 1320
aaaaaaaaa a 1331

```

<210> 8

<211> 349

<212> PRT

<213> Glycine max

<400> 8

```

Met Ala Thr Val Val Arg Ile Pro Thr Ile Ser Cys Ile His Ile His
  1             5             10             15

Thr Phe Arg Ser Gln Ser Pro Arg Thr Phe Ala Arg Ile Arg Val Gly
          20             25             30

Pro Arg Ser Trp Ala Pro Ile Arg Ala Ser Ala Ala Ser Ser Glu Arg
          35             40             45

Gly Glu Ile Val Leu Glu Gln Lys Pro Lys Lys Asp Asp Lys Lys Lys
          50             55             60

Leu Gln Lys Gly Ile Ala Glu Phe Tyr Asp Glu Ser Ser Gly Leu Trp
          65             70             75             80

Glu Asn Ile Trp Gly Asp His Met His His Gly Phe Tyr Asp Ser Asp
          85             90             95

Ser Thr Val Ser Leu Ser Asp His Arg Ala Ala Gln Ile Arg Met Ile
          100            105            110

Gln Glu Ser Leu Arg Phe Ala Ser Val Ser Glu Glu Arg Ser Lys Trp
          115            120            125

Pro Lys Ser Ile Val Asp Val Gly Cys Gly Ile Gly Gly Ser Ser Arg
          130            135            140

Tyr Leu Ala Lys Lys Phe Gly Ala Thr Ser Val Gly Ile Thr Leu Ser
          145            150            155            160

```

Pro Val Gln Ala Gln Arg Ala Asn Ala Leu Ala Ala Ala Gln Gly Leu
 165 170 175

Ala Asp Lys Val Ser Phe Gln Val Ala Asp Ala Leu Gln Gln Pro Phe
 180 185 190

Ser Asp Gly Gln Phe Asp Leu Val Trp Ser Met Glu Ser Gly Glu His
 195 200 205

Met Pro Asp Lys Ala Lys Phe Val Gly Glu Leu Ala Arg Val Ala Ala
 210 215 220

Pro Gly Ala Ile Ile Ile Ile Val Thr Trp Cys His Arg Asp Leu Gly
 225 230 235 240

Pro Asp Glu Gln Ser Leu His Pro Trp Glu Gln Asp Leu Leu Lys Lys
 245 250 255

Ile Cys Asp Ala Tyr Tyr Leu Pro Ala Trp Cys Ser Thr Ser Asp Tyr
 260 265 270

Val Lys Leu Leu Gln Ser Leu Ser Leu Gln Asp Ile Lys Ser Glu Asp
 275 280 285

Trp Ser Arg Phe Val Ala Pro Phe Trp Pro Ala Val Ile Arg Ser Ala
 290 295 300

Phe Thr Trp Lys Gly Leu Ser Ser Leu Leu Ser Ser Gly Lys Leu Gly
 305 310 315 320

Ile Tyr Ile Ala Phe Gln Lys Gln Thr Pro Pro Ser Ser Ile Ala Thr
 325 330 335

Cys Lys Ser Tyr Val Thr Asp His Tyr Phe His Thr Arg
 340 345

<210> 9
 <211> 1011
 <212> DNA
 <213> Triticum aestivum

<400> 9
 acgagtcgtc cggcctgtgg gagagcatct ggggcgagca catgcaccac ggcttctacg 60
 actccggcga ggccgcctcc atgtccgacc accgccgcgc ccagatccgc atgatcgagg 120
 aggcctctgc cttcgccgcc gtccccgacg atccgacaaa caaacccaaa acgattgttg 180
 atgttggatg cggaaatcggg ggtagctcaa gatacctggg cgaacaaata tggagcacia 240
 tgctctggga tcacattgac ccagtgcaag ctgagagagg aaatgccctc gcggcagcgc 300
 aagggttgtt cgggacaagg ttctttccaa ttgctgatct ctgggagcaa ccatttcctg 360
 gatgggcatt tgatcttgtc cgggnccatg ggagantggt gacacatgcc gaacaaacag 420
 aagtttgtaa gcgagctggc acgcgtcgca gctccaggag caactatcat catcgtgacc 480
 tgggtgccata ggaacctcgc gccatcggag gactcactga aacctgacga gctgaatctt 540
 ttgaaaaaga tttgtgatgc atattacctc ccggattggt gctctccctc ggattatgtc 600
 aagattgccg agtcattgtc tcttgaggat atcaaaacgg ccgactggtc tgaaaacgtg 660
 gccccgttct ggctgtctgt catccaatca gcactgacat ggaaaggcct cacttctcta 720
 ctaaggagtg gatggaagac gataaaggga gcactggtga tgctctcat gatccaaggc 780
 tacaagaaag gcctcattaa gttcaagcat catcacctgc cacaaccccc aagcagccat 840
 agaaggagaa cctggaggcc gcacgcacca agagtggtag aatagaacca tgtgattgga 900
 atagactcng cttgctgtcg ccttggtagc tgaataatc gtgttaccgt gcctctgtat 960
 ctgcaactgg aagtgccata tgagaatggt tcctaaaagc aaaatctcct c 1011

<210> 10
 <211> 293
 <212> PRT
 <213> Triticum aestivum

 <400> 10
 Glu Ser Ser Gly Leu Trp Glu Ser Ile Trp Gly Glu His Met His His
 1 5 10 15
 Gly Phe Tyr Asp Ser Gly Glu Ala Ala Ser Met Ser Asp His Arg Arg
 20 25 30
 Ala Gln Ile Arg Met Ile Glu Glu Ala Leu Ala Phe Ala Ala Val Pro
 35 40 45
 Asp Asp Pro Thr Asn Lys Pro Lys Thr Ile Val Asp Val Gly Cys Gly
 50 55 60
 Ile Gly Gly Ser Ser Arg Tyr Leu Gly Glu Gln Ile Trp Ser Thr Met
 65 70 75 80
 Leu Trp Asp His Ile Asp Pro Val Gln Ala Glu Arg Gly Asn Ala Leu
 85 90 95
 Ala Ala Ala Gln Gly Val Val Arg Thr Arg Phe Phe Pro Ile Ala Asp
 100 105 110
 Leu Trp Glu Gln Pro Phe Pro Gly Trp Ala Phe Asp Leu Val Xaa Xaa
 115 120 125
 Xaa Xaa Xaa Xaa Xaa His Met Pro Asn Lys Gln Lys Phe Val Ser Glu
 130 135 140
 Leu Ala Arg Val Ala Ala Pro Gly Ala Thr Ile Ile Ile Val Thr Trp
 145 150 155 160
 Cys His Arg Asn Leu Ala Pro Ser Glu Asp Ser Leu Lys Pro Asp Glu
 165 170 175
 Leu Asn Leu Leu Lys Lys Ile Cys Asp Ala Tyr Tyr Leu Pro Asp Trp
 180 185 190
 Cys Ser Pro Ser Asp Tyr Val Lys Ile Ala Glu Ser Leu Ser Leu Glu
 195 200 205
 Asp Ile Lys Thr Ala Asp Trp Ser Glu Asn Val Ala Pro Phe Trp Pro
 210 215 220
 Ala Val Ile Gln Ser Ala Leu Thr Trp Lys Gly Leu Thr Ser Leu Leu
 225 230 235 240
 Arg Ser Gly Trp Lys Thr Ile Lys Gly Ala Leu Val Met Pro Leu Met
 245 250 255
 Ile Gln Gly Tyr Lys Lys Gly Leu Ile Lys Phe Lys His His His Leu
 260 265 270
 Pro Gln Thr Pro Ser Ser His Arg Arg Arg Thr Trp Arg Pro His Arg
 275 280 285

Pro Arg Val Val Glu
290

<210> 11
<211> 432
<212> DNA
<213> Oryza sativa

<400> 11
ccacgtcgag ctctggtgag ccgacgcgc gtcgcgcgc ggccggttcg ccttcgcct 60
ggcgcgccg ctgcgcgcca ggtccgacct ctccacggg aactccgcgc acgcctcct 120
cctcctccgc tccgcctcgc tcgcgttcct cttcacgcgc ccctacggcg gcgaccacgg 180
cgtcggcgcg gacgcggcca ccaccgcctc catcccttc ttctccctt cctttctccc 240
cgctcctgga tcaggccaca ggaggggagc gatggtggag gcggccaccg taggcggcg 300
aggtggcgtt cctcctcct agctcccaga cccggctgga ggagggagtg atggtggcg 360
nagggcgngc tccctcctct ccctccctcc tcctcacaat ntggccggag ggaggaang 420
gccgcggncc aa 432

<210> 12
<211> 75
<212> PRT
<213> Oryza sativa

<400> 12
His Val Glu Leu Trp Cys Ala Asp Ala Ala Ser Ala Ala Gly Arg Phe
1 5 10 15
Ala Phe Ala Leu Gly Ala Pro Leu Ala Ala Arg Ser Asp Leu Ser Thr
20 25 30
Gly Asn Ser Ala His Ala Ser Leu Leu Leu Arg Ser Ala Ser Val Ala
35 40 45
Phe Leu Phe Thr Ala Pro Tyr Gly Gly Asp His Gly Val Gly Ala Asp
50 55 60
Ala Ala Thr Thr Ala Ser Ile Pro Ser Phe Ser
65 70 75

<210> 13
<211> 628
<212> DNA
<213> Oryza sativa

<400> 13
gaagagctac ggcctccgcc ggttcgacca cgtcgtcggc aacgtgccgg agctcgtcc 60
ggtageccgc tacatctccg ggttcaccgg gtccacgag ttccgccagt tcaccgccga 120
ggacgtgggc accgccgaga gcggcctcaa ctccggtggtg ctccccaaca acgcggagac 180
cgtgctgctg ccgctcaacg agccggtgca cggcaccaag cggcgagacc agatacagac 240
gtacctggac caccacggcg gcccgggggt gcagcacatc gcgctggcca gcgacgacgt 300
gctcgggacg ctganggaga tgcnggcgc ctccgcatgg gcggttcgat tcttgggccc 360
gccgcgcgca actactacga cggctgcgcg gcgcncggg acttctctcg ggagagcaat 420
taacaatgcc aagactcngg tgtcctggac aaggatacaa gggtttcaa tnttaacaag 480
cattgaanag nnactttctg gngagatcaa gatggtgatg aaagtnaatg gaagntncaa 540
aggggntcgc ggttggaaga atntcggtt aatcataggg tngaaacctn agcacagcct 600
anttaggtca gagatngcn ganaaatt 628

<210> 14
<211> 123

<212> PRT

<213> Oryza sativa

<400> 14

Tyr Gly Leu Arg Arg Phe Asp His Val Val Gly Asn Val Pro Glu Leu
 1 5 10 15

Ala Pro Val Ala Ala Tyr Ile Ser Gly Phe Thr Gly Phe His Glu Phe
 20 25 30

Ala Glu Phe Thr Ala Glu Asp Val Gly Thr Ala Glu Ser Gly Leu Asn
 35 40 45

Ser Val Val Leu Ala Asn Asn Ala Glu Thr Val Leu Leu Pro Leu Asn
 50 55 60

Glu Pro Val His Gly Thr Lys Arg Arg Ser Gln Ile Gln Thr Tyr Leu
 65 70 75 80

Asp His His Gly Gly Pro Gly Val Gln His Ile Ala Leu Ala Ser Asp
 85 90 95

Asp Val Leu Gly Thr Leu Xaa Glu Met Pro Gly Ala Ser Ala Trp Ala
 100 105 110

Val Arg Phe Leu Gly Pro Pro Pro Pro Thr Thr
 115 120

<210> 15

<211> 1027

<212> DNA

<213> Glycine max

<400> 15

tcacaccaca ccaatgccaa taccatgtg caacgaaatt caagcccaag cccaagccca 60
 agcccaagcc caacctgggt ttaagctcgt cggtttcaaa aacttcgtcc gaaccaatcc 120
 taagtccggac cgctttcaag tcaaccgctt ccaccacatc gagttctggt gcaccgatgc 180
 caccaacgcc tctcgccgat tctcttgagg acttggaatg cctattgtgg caaaatctga 240
 tctctccacc ggaacccaaa tccacgcctc ctacctcctc cgctccggcg acctctcctt 300
 cctcttctcc gctccttact ctcctctctc ctccgcggcg tctccgctg cctcctccgc 360
 ctccattccc agtttcgacg ccgccacctg ccttgccctc gctgccaaac acggcttcgg 420
 cgtccgcgcc atcgcttggt aagtcgccga cgcggaagcc gctttcagcg ccagcgtcgc 480
 gaaaggagcc gagccggcgt cgcgcgggt tctcgtcgac gatcgaccg gcttcgcgga 540
 ggtgcgcctc tacggcgacg tgggtgctccg ctacgtcagc tacaaggacg ccgcgccgca 600
 ggcgccacac gcagatncgt cgcggtggtt cctgccggga ttcgaggccg cggcgctcgc 660
 gtcttcggtt ccggagctgg actacgggat ccggcggtg gaccacgccg tcgggaacgt 720
 tccggagctg gcgccggcg tgaggtacct gaaaggcttc agcggattcc acgagttcgc 780
 ggagttcacc gcggaggacg tgggaacgag cgagagcggg ttgaactcng tggttctggc 840
 ngaacaactc ggagacggtg ttgctgccgc tgaacnagcc cggtttacgg aacgaaagag 900
 gaagaagcca nattgagnnc gtatttngaa cacaancnaa aggtgcttgg tgtgcagcaa 960
 ccttgcgcnt tgttactcnc naacatctc ancacactga ggagagatg anaaaanccg 1020
 acgtttg 1027

<210> 16

<211> 276

<212> PRT

<213> Glycine max

<400> 16

Met Pro Ile Pro Met Cys Asn Glu Ile Gln Ala Gln Ala Gln Ala Gln
 1 5 10 15

Ala Gln Ala Gln Pro Gly Phe Lys Leu Val Gly Phe Lys Asn Phe Val
 20 25 30

Arg Thr Asn Pro Lys Ser Asp Arg Phe Gln Val Asn Arg Phe His His
 35 40 45

Ile Glu Phe Trp Cys Thr Asp Ala Thr Asn Ala Ser Arg Arg Phe Ser
 50 55 60

Trp Gly Leu Gly Met Pro Ile Val Ala Lys Ser Asp Leu Ser Thr Gly
 65 70 75 80

Asn Gln Ile His Ala Ser Tyr Leu Leu Arg Ser Gly Asp Leu Ser Phe
 85 90 95

Leu Phe Ser Ala Pro Tyr Ser Pro Ser Leu Ser Ala Gly Ser Ser Ala
 100 105 110

Ala Ser Ser Ala Ser Ile Pro Ser Phe Asp Ala Ala Thr Cys Leu Ala
 115 120 125

Phe Ala Ala Lys His Gly Phe Gly Val Arg Ala Ile Ala Leu Glu Val
 130 135 140

Ala Asp Ala Glu Ala Ala Phe Ser Ala Ser Val Ala Lys Gly Ala Glu
 145 150 155 160

Pro Ala Ser Pro Pro Val Leu Val Asp Asp Arg Thr Gly Phe Ala Glu
 165 170 175

Val Arg Leu Tyr Gly Asp Val Val Leu Arg Tyr Val Ser Tyr Lys Asp
 180 185 190

Ala Ala Pro Gln Ala Pro His Ala Asp Xaa Ser Arg Trp Phe Leu Pro
 195 200 205

Gly Phe Glu Ala Ala Ala Ser Ser Ser Ser Phe Pro Glu Leu Asp Tyr
 210 215 220

Gly Ile Arg Arg Leu Asp His Ala Val Gly Asn Val Pro Glu Leu Ala
 225 230 235 240

Pro Ala Val Arg Tyr Leu Lys Gly Phe Ser Gly Phe His Glu Phe Ala
 245 250 255

Glu Phe Thr Ala Glu Asp Val Gly Thr Ser Glu Ser Gly Leu Asn Ser
 260 265 270

Val Val Leu Ala
 275

<210> 17

<211> 511

<212> DNA

<213> Vernonia mesipifolia

<400> 17
ccacaccgat tgccggaact tcaccgcctc tcacggcctt gcagtcgag caatcgccat 60
tgaagtcgat gacgccgaat tagctttctc cgtcagcgtc tctcacggcg ctaaaccctc 120
cgctgctcct gtaacccttg gaaacaacga cgtcgtattg tctgaagtta agctttacgg 180
cgatgtcgct ttccggtaca taagttacaa aaatccgaac tatacatctt cctttttgcc 240
cgggttcgag cccgttgaaa agacgtcgtc gttttatgac cttgactacg gtatccgccg 300
tttgaccac gccgtaggaa cgtccctgag cttgcttcgg cagtggacta cgtgaaatca 360
ttcaccggat tccatgagtt cgccgaattc accgcggagg acgtcgggac gacgagagg 420
gaactgaatt cggtcgtttt agcttgcaac agtgagatgg tcttgattcc gatgaacgag 480
ccggtgtacg gaanaaaagg aagagccaga t 511

<210> 18
<211> 170
<212> PRT
<213> *Vernonia mesipifolia*

<400> 18
His Thr Asp Cys Arg Asn Phe Thr Ala Ser His Gly Leu Ala Val Arg
1 5 10 15
Ala Ile Ala Ile Glu Val Asp Asp Ala Glu Leu Ala Phe Ser Val Ser
20 25 30
Val Ser His Gly Ala Lys Pro Ser Ala Ala Pro Val Thr Leu Gly Asn
35 40 45
Asn Asp Val Val Leu Ser Glu Val Lys Leu Tyr Gly Asp Val Ala Phe
50 55 60
Arg Tyr Ile Ser Tyr Lys Asn Pro Asn Tyr Thr Ser Ser Phe Leu Pro
65 70 75 80
Gly Phe Glu Pro Val Glu Lys Thr Ser Ser Phe Tyr Asp Leu Asp Tyr
85 90 95
Gly Ile Arg Arg Leu Asp His Ala Val Gly Asn Val Pro Glu Leu Ala
100 105 110
Ser Ala Val Asp Tyr Val Lys Ser Phe Thr Gly Phe His Glu Phe Ala
115 120 125
Glu Phe Thr Ala Glu Asp Val Gly Thr Ser Glu Arg Glu Leu Asn Ser
130 135 140
Val Val Leu Ala Cys Asn Ser Glu Met Val Leu Ile Pro Met Asn Glu
145 150 155 160
Pro Val Tyr Gly Xaa Lys Gly Arg Ala Arg
165 170

<210> 19
<211> 1165
<212> DNA
<213> *Triticum aestivum*

<400> 19
caagaagcga acacacacca tgccgcccac ccccaccacc cccgcagcca ccggcgccgc 60
cgcggtgacg ccggagcacg cgcgccgcg cgaatggtc cgcttcaacc ccgcgacgca 120
ccgcttcac acgtcgcct tccaccacgt cgagttctgg tgcgcggacg ccgcctccgc 180
cgccggccgc ttcgcttcg cgctcggcgc gccctcgc gccaggtccg acctctccac 240

```

ggggaactcc gtgcacgect cccagctgct ccgctcgggc aacctcgect tcctcttcac 300
cgcgccctac gccaacggct ggcacgccgc caccgcctcc ctgccctect tctccgccga 360
cgccgcgcgc cggttctccg cggaccacgg gctcgcggtg cgctccatag cgctgcgcgt 420
cgcgagacgc gccgaggect tccgcgccag cgtcgacggg ggcgcgcgcc cggccttcag 480
ccccgtggac ctcggccgcg gtttcggctt tgcggaggtc gagctctacg ggcacgtcgt 540
gctccgcttc gtcagcatcc ggacggnacg gacgtgcctt ctgcccggg ttcgangggc 600
ttgagcaacc gggtgccgtg gactaanggc tgacacggnt tgacacgttg tccgnaagtc 660
cggagcttgc ttcggcgccg cctaactag cgggctnaac gggttcaana attcgccagt 720
taacacggag gacgtgggca cggccgagag cgggctcaac tcgatggtgc tcgccaacaa 780
ctcggagggc gtgctgctgc cgctcaacga gccggtgcac ggcaccaagc gccggagcca 840
gatacagacg ttcctggaac accacggcgg ctcgggcggtg cagcacatcg cggtggccag 900
cagcgacgtg ctcaggacgc tcaggagat gcgtgcgcgc tccgccatgg gcggttcga 960
cttctgcca ccccgctgc cgaagtacta cgaaggcgtg cggcgcatcg ccggggatgt 1020
gctctcggag gcgcaaatna aggaatgcaa gaactggggg tgctcntcca caaggaagaa 1080
caaagggtgt tgctacaaat cctcaacaag ccaatntggg acaagccgac ttgttctcgt 1140
agatattcac angatctggt gcatg 1165

```

<210> 20
 <211> 179
 <212> PRT
 <213> *Triticum aestivum*

<400> 20
 Met Pro Pro Thr Pro Thr Thr Pro Ala Ala Thr Gly Ala Ala Ala Val
 1 5 10 15
 Thr Pro Glu His Ala Arg Pro Arg Arg Met Val Arg Phe Asn Pro Arg
 20 25 30
 Ser Asp Arg Phe His Thr Leu Ala Phe His His Val Glu Phe Trp Cys
 35 40 45
 Ala Asp Ala Ala Ser Ala Ala Gly Arg Phe Ala Phe Ala Leu Gly Ala
 50 55 60
 Pro Leu Ala Ala Arg Ser Asp Leu Ser Thr Gly Asn Ser Val His Ala
 65 70 75 80
 Ser Gln Leu Leu Arg Ser Gly Asn Leu Ala Phe Leu Phe Thr Ala Pro
 85 90 95
 Tyr Ala Asn Gly Cys Asp Ala Ala Thr Ala Ser Leu Pro Ser Phe Ser
 100 105 110
 Ala Asp Ala Ala Arg Arg Phe Ser Ala Asp His Gly Leu Ala Val Arg
 115 120 125
 Ser Ile Ala Leu Arg Val Ala Asp Ala Ala Glu Ala Phe Arg Ala Ser
 130 135 140
 Val Asp Gly Gly Ala Arg Pro Ala Phe Ser Pro Val Asp Leu Gly Arg
 145 150 155 160
 Gly Phe Gly Phe Ala Glu Val Glu Leu Tyr Gly Asp Val Val Leu Arg
 165 170 175
 Phe Val Ser

<210> 21
 <211> 1102
 <212> DNA
 <213> Zea mays

<220>
 <221> unsure
 <222> (454)

<220>
 <221> unsure
 <222> (1072)

<220>
 <221> unsure
 <222> (1083)

<220>
 <221> unsure
 <222> (1092)

<220>
 <221> unsure
 <222> (1100)

<400> 21
 atggctcacg cggcgctgct ccattgctcc cagtcctcca ggagcctcgc agcctgccgc 60
 cgcggcagyc actaccgcgc cccttcgcac gtcccgcgcc actcccgccg tctccgacgc 120
 gccgtcgtca gcctgcgtcc gatggcctcg tcgacggctc aggcccccgc gacggcgccg 180
 ccgggtctga aggaggcat cgcggggctg tacgacgagt cgtcggggct gtgggagaac 240
 atctggggcg accacatgca ccacggcttc tacgactcga gcgaggccgc ctccatggcc 300
 gatcaccgcc gcgccagat ccgcatgac gaggaggcgc tcgccttcgc cgggtgtcca 360
 gcctcagatg atccagagaa gacacaaaa acaatagtcg atgtcggatg tggcattggt 420
 ggtagctcaa ggtacttggc gaagaaatac ggancgcagt gcactgggat cacgttgagc 480
 cctgttcaag ccgagagagg aaatgctctc gctgcagcgc aggggttgtc ggatcagggt 540
 actctgcaag ttgctgatgc tctggagcaa ccgtttcctg acgggcagtt cgatctggtg 600
 tgggtccatgg agagtggcga gcacatgccg gacaagagaa agtttgtag tgagctagca 660
 cgcgtggcgg ctccctggagg gacaataatc atcgtgacat ggtgccatag gaacctggat 720
 ccatccgaaa cctcgctaaa gcccgatgaa ctgagcctcc tgaggaggat atgcgacgcg 780
 tactacctcc cggactggtg ctacacctca gactatgtga acattgccaa gtcactgtct 840
 ctcgaggata tcaagacagc tgactggctg gagaacgtgg ccccgttttg gcccgccgtg 900
 ataaaaatcag cgctaacatg gaagggttc acctctctgc tgacgaccgg atggaagacg 960
 atcagaggcg cgatggtgat gccgctaag atccagggt acaagaagg gctcatcaaa 1020
 ttcacatca tcacctgtcg caagcctgga gccgcgtagt gatctatacc gnccacggcg 1080
 tenttaactc tnacggaaan ct 1102

<210> 22
 <211> 352
 <212> PRT
 <213> Zea mays

<220>
 <221> UNSURE
 <222> (152)

<400> 22
 Met Ala His Ala Ala Leu Leu His Cys Ser Gln Ser Ser Arg Ser Leu
 1 5 10 15

Ala Ala Cys Arg Arg Gly Ser His Tyr Arg Ala Pro Ser His Val Pro
 20 25 30
 Arg His Ser Arg Arg Leu Arg Arg Ala Val Val Ser Leu Arg Pro Met
 35 40 45
 Ala Ser Ser Thr Ala Gln Ala Pro Ala Thr Ala Pro Pro Gly Leu Lys
 50 55 60
 Glu Gly Ile Ala Gly Leu Tyr Asp Glu Ser Ser Gly Leu Trp Glu Asn
 65 70 75 80
 Ile Trp Gly Asp His Met His His Gly Phe Tyr Asp Ser Ser Glu Ala
 85 90 95
 Ala Ser Met Ala Asp His Arg Arg Ala Gln Ile Arg Met Ile Glu Glu
 100 105 110
 Ala Leu Ala Phe Ala Gly Val Pro Ala Ser Asp Asp Pro Glu Lys Thr
 115 120 125
 Pro Lys Thr Ile Val Asp Val Gly Cys Gly Ile Gly Gly Ser Ser Arg
 130 135 140
 Tyr Leu Ala Lys Lys Tyr Gly Xaa Gln Cys Thr Gly Ile Thr Leu Ser
 145 150 155 160
 Pro Val Gln Ala Glu Arg Gly Asn Ala Leu Ala Ala Ala Gln Gly Leu
 165 170 175
 Ser Asp Gln Val Thr Leu Gln Val Ala Asp Ala Leu Glu Gln Pro Phe
 180 185 190
 Pro Asp Gly Gln Phe Asp Leu Val Trp Ser Met Glu Ser Gly Glu His
 195 200 205
 Met Pro Asp Lys Arg Lys Phe Val Ser Glu Leu Ala Arg Val Ala Ala
 210 215 220
 Pro Gly Gly Thr Ile Ile Ile Val Thr Trp Cys His Arg Asn Leu Asp
 225 230 235 240
 Pro Ser Glu Thr Ser Leu Lys Pro Asp Glu Leu Ser Leu Leu Arg Arg
 245 250 255
 Ile Cys Asp Ala Tyr Tyr Leu Pro Asp Trp Cys Ser Pro Ser Asp Tyr
 260 265 270
 Val Asn Ile Ala Lys Ser Leu Ser Leu Glu Asp Ile Lys Thr Ala Asp
 275 280 285
 Trp Ser Glu Asn Val Ala Pro Phe Trp Pro Ala Val Ile Lys Ser Ala
 290 295 300
 Leu Thr Trp Lys Gly Phe Thr Ser Leu Leu Thr Thr Gly Trp Lys Thr
 305 310 315 320
 Ile Arg Gly Ala Met Val Met Pro Leu Met Ile Gln Gly Tyr Lys Lys
 325 330 335

Gly Leu Ile Lys Phe Thr Ile Ile Thr Cys Arg Lys Pro Gly Ala Ala
340 345 350

<210> 23
<211> 521
<212> DNA
<213> Oryza sativa

<220>
<221> unsure
<222> (269)

<220>
<221> unsure
<222> (274)

<220>
<221> unsure
<222> (286)

<220>
<221> unsure
<222> (302)

<220>
<221> unsure
<222> (330)

<220>
<221> unsure
<222> (381)..(382)

<220>
<221> unsure
<222> (387)

<220>
<221> unsure
<222> (398)

<220>
<221> unsure
<222> (418)

<220>
<221> unsure
<222> (429)

<220>
<221> unsure
<222> (436)

<220>
<221> unsure
<222> (462)

<220>
<221> unsure
<222> (467)

<220>

<221> unsure

<222> (473)

<220>

<221> unsure

<222> (514)

<400> 23

```

cttacagaca aacggcagtt tgtaagcgag ctggcacgcg tcgcagctcc tggggcgaga 60
ataatcattg tgacctggtg ccataggaac ctcgagccat ccgaagagtc cctgaaacct 120
gatgagctga atctcctgaa aaggatatgc gatgcatatt atctcccaga ctggtgctct 180
ccttctgatt atgtcaaaat tgccgagtcg ctgtctcttg aggatataag gacagctgat 240
tggtcaagag aacgtcgccc caatccggnc tgcnggttat taaatnaagc aattgacatg 300
gnaagggtta actttctcct ggctaagaan tgggtgggaa gacgattaag aaggtggaat 360
gggtgatgcc tccggatgat nnaaggntac aaagaaangg gtcaacaaat ttaacaanaa 420
caacctgtnc caaagncccg aaacaacgca ataatacccc antaatnaaa ttncgctcct 480
ggctaacctt ctccaacaac gaattaatgg aaanttctga c 521

```

<210> 24

<211> 172

<212> PRT

<213> Oryza sativa

<400> 24

```

Phe Arg His Gly His Ala Leu Ala Gln Pro Phe Pro Asp Gly Gln Phe
  1             5             10             15

```

```

Asp Leu Val Trp Ser Met Glu Ser Asp Glu His Met Pro Asp Lys Arg
      20             25             30

```

```

Gln Phe Val Ser Glu Leu Ala Arg Val Ala Ala Pro Gly Ala Arg Ile
      35             40             45

```

```

Ile Ile Val Thr Trp Cys His Arg Asn Leu Glu Pro Ser Glu Glu Ser
      50             55             60

```

```

Leu Lys Pro Asp Glu Leu Asn Leu Leu Lys Arg Ile Cys Asp Ala Tyr
      65             70             75             80

```

```

Tyr Leu Pro Asp Trp Cys Ser Pro Ser Asp Tyr Val Lys Ile Ala Glu
      85             90             95

```

```

Ser Leu Ser Leu Glu Asp Ile Arg Thr Ala Asp Trp Ser Glu Asn Val
      100            105            110

```

```

Ala Pro Phe Trp Pro Ala Val Ile Lys Ser Ala Leu Thr Trp Lys Gly
      115            120            125

```

```

Leu Thr Ser Leu Leu Arg Ser Gly Trp Glu Thr Val Arg Gly Ala Met
      130            135            140

```

```

Val Met Pro Leu Val Ile Glu Gly Tyr Lys Lys Gly Leu Ile Lys Phe
      145            150            155            160

```

```

Pro Ile Ile Thr Cys Arg Lys Pro Glu Thr Thr Gln
      165            170

```

<210> 25

<211> 464

<212> DNA
<213> *Oryza sativa*

<400> 25
gcacgagtag agccacggg cgaactggca cgcgtgcac cactgctccg ctgcacgagc 60
cgtcatctct gcgcctcggc ttccctcgc gccggcctct gcctccacca ccaccgccgc 120
cgccgccgca gcagccggag gacgaaactc gccgtgcgcg cgatggcacc gacgttgctc 180
tcgtcgtcga cggcggcggc agctcccccg gggctgaagg agggcatcgc ggggtcttac 240
gacgagtcgt ccggcgtgtg ggagagcatc tggggcgagc acatgcacca cggcttctac 300
gacgccggcg aggccgcctc catgtccgac caccgccgcg cccagatccg catgatcgag 360
gaatccctcg ccttcgccgc cgttccccga tgatgcgggt aacaaaccca aaagtgttat 420
ttactgtttg gtgttgcaaa tgggggtacc tccaaaaaac ttg 464

<210> 26
<211> 128
<212> PRT
<213> *Oryza sativa*

<400> 26
Ala Arg Val Gln Pro Thr Gly Ala Leu Ala Pro Leu His Pro Leu Leu
1 5 10 15
Arg Cys Thr Ser Arg His Leu Cys Ala Ser Ala Ser Pro Arg Ala Gly
20 25 30
Leu Cys Leu His His His Arg Arg Arg Arg Arg Ser Ser Arg Arg Thr
35 40 45
Lys Leu Ala Val Arg Ala Met Ala Pro Thr Leu Ser Ser Ser Ser Thr
50 55 60
Ala Ala Ala Ala Pro Pro Gly Leu Lys Glu Gly Ile Ala Gly Leu Tyr
65 70 75 80
Asp Glu Ser Ser Gly Val Trp Glu Ser Ile Trp Gly Glu His Met His
85 90 95
His Gly Phe Tyr Asp Ala Gly Glu Ala Ala Ser Met Ser Asp His Arg
100 105 110
Arg Ala Gln Ile Arg Met Ile Glu Glu Ser Leu Ala Phe Ala Ala Val
115 120 125

<210> 27
<211> 1189
<212> DNA
<213> Glycine max

<400> 27
ggacatggcc accgtggtga ggatcccaac aatctcatgc atccacatcc acacgttccg 60
ttcccaatcc cctcgactt tcgccagaat ccgggtcggc cccaggtcgt gggctcctat 120
tcgggcatcg gcagcgagct cggagagagg ggagatagta ttggagcaga agccgaagaa 180
ggatgacaag aagaagctgc agaagggaat cgcagagttt tacgacgagt cgtctggctt 240
atgggagaac atttggggcg accacatgca ccatggcttt tatgactcgg attccactgt 300
ttcgctttcg gatcatcgtg ctgctcagat ccgaatgatc caagagtctc ttcgctttgc 360
ctctgtttct gaggagcgtg gtaaatggcc caagagtata gttgatgttg ggtgtggcat 420
aggtggcagc tctagatacc tggccaagaa atttggagca accagtgtag gcatcactct 480
gagtcctgtt caagctcaaa gagcaaatgc tcttgctgct gctcaaggat tggctgataa 540
ggtttccttt caggttgctg acgctctaca gcaaccattc tctgacggcc agtttgatct 600
ggtgtggtcc atggagagtg gagagcatat gcctgacaaa gctaagtttg ttggagagtt 660

```

agctcgggta gcagcaccag gtgccattat aataatagta acatgggtgcc acagggatct 720
tgcccttgac gaacaatcct tacatccatg ggagcaagat ctcttaaaga agatttgca 780
tgcataattac ctccctgcct ggtgctcaac ttctgattat gttaagttgc tccaatccct 840
gtcacttcag gacatcaagt cagaagattg gtctcgcttt gttgctccat tttggccagc 900
agtgatacgc tcagccttca catggaaggg tctatcttca ctcttgagca gtggacaaaa 960
aacgataaaa ggagcttttg ctatgccatt gatgatagag ggatacaaga aagatctaata 1020
taagtttgcc atcattacat gtcgaaaacc tgaataaatg gagaggcagg attactttta 1080
tagaatgaac caagtttcca acaggtcggt tatttcgata gttgagaaac aagagaaaaa 1140
ataaatgaaa ggggttggtc gattttaaaa aaaaaaaaaa aaaaaaaaaa 1189

```

<210> 28
 <211> 350
 <212> PRT
 <213> Glycine max

<400> 28
 Met Ala Thr Val Val Arg Ile Pro Thr Ile Ser Cys Ile His Ile His
 1 5 10 15
 Thr Phe Arg Ser Gln Ser Pro Arg Thr Phe Ala Arg Ile Arg Val Gly
 20 25 30
 Pro Arg Ser Trp Ala Pro Ile Arg Ala Ser Ala Ala Ser Ser Glu Arg
 35 40 45
 Gly Glu Ile Val Leu Glu Gln Lys Pro Lys Lys Asp Asp Lys Lys Lys
 50 55 60
 Leu Gln Lys Gly Ile Ala Glu Phe Tyr Asp Glu Ser Ser Gly Leu Trp
 65 70 75 80
 Glu Asn Ile Trp Gly Asp His Met His His Gly Phe Tyr Asp Ser Asp
 85 90 95
 Ser Thr Val Ser Leu Ser Asp His Arg Ala Ala Gln Ile Arg Met Ile
 100 105 110
 Gln Glu Ser Leu Arg Phe Ala Ser Val Ser Glu Glu Arg Ser Lys Trp
 115 120 125
 Pro Lys Ser Ile Val Asp Val Gly Cys Gly Ile Gly Gly Ser Ser Arg
 130 135 140
 Tyr Leu Ala Lys Lys Phe Gly Ala Thr Ser Val Gly Ile Thr Leu Ser
 145 150 155 160
 Pro Val Gln Ala Gln Arg Ala Asn Ala Leu Ala Ala Ala Gln Gly Leu
 165 170 175
 Ala Asp Lys Val Ser Phe Gln Val Ala Asp Ala Leu Gln Gln Pro Phe
 180 185 190
 Ser Asp Gly Gln Phe Asp Leu Val Trp Ser Met Glu Ser Gly Glu His
 195 200 205
 Met Pro Asp Lys Ala Lys Phe Val Gly Glu Leu Ala Arg Val Ala Ala
 210 215 220
 Pro Gly Ala Ile Ile Ile Ile Val Thr Trp Cys His Arg Asp Leu Gly
 225 230 235 240

Pro Asp Glu Gln Ser Leu His Pro Trp Glu Gln Asp Leu Leu Lys Lys
 245 250 255

Ile Cys Asp Ala Tyr Tyr Leu Pro Ala Trp Cys Ser Thr Ser Asp Tyr
 260 265 270

Val Lys Leu Leu Gln Ser Leu Ser Leu Gln Asp Ile Lys Ser Glu Asp
 275 280 285

Trp Ser Arg Phe Val Ala Pro Phe Trp Pro Ala Val Ile Arg Ser Ala
 290 295 300

Phe Thr Trp Lys Gly Leu Ser Ser Leu Leu Ser Ser Gly Gln Lys Thr
 305 310 315 320

Ile Lys Gly Ala Leu Ala Met Pro Leu Met Ile Glu Gly Tyr Lys Lys
 325 330 335

Asp Leu Ile Lys Phe Ala Ile Ile Thr Cys Arg Lys Pro Glu
 340 345 350

<210> 29
 <211> 1257
 <212> DNA
 <213> Triticum aestivum

<220>
 <221> unsure
 <222> (31)

<220>
 <221> unsure
 <222> (151)

<400> 29
 gaggtctccaa atacaaaatg gcaaactccg nccgcccctgc tccactcact cctctccacc 60
 gcctggagcgc cgcgcgcgcg cctcgaccga gcctcggcca cgcggctcgc cccgtccccc 120
 ggctgtctct ggcgtctctc ccgcccagac ngctccgtgc gcccgatggc gtcgtcgacg 180
 accgcggccc gggcgacgcg gcgcgcgcgc ggctgaagga gggcatcgcg gggctctacg 240
 acgagtcgtc cggcctgtgg gagagcatct ggggcgagca catgcaccac ggcttctacg 300
 actccggcga ggccgcctcc atgtccgacc accgcgcgcg ccagatccgc atgatcgagg 360
 aggcctcgcg cttcgccgcg gtcccgcgac atccgacaaa caaacccaaa acgattgttg 420
 atgttggtatg cggaatcggg ggtagctcaa gatacctggc gaacaaatat ggagcacaat 480
 gctctgggat cacattgagc ccagtgcagg ctgagagagg aaatgccctc gcggcagcgc 540
 aggggttgtc ggacaaggct tctttccaag ttgctgatgc tctggagcaa ccatttcctg 600
 atgggcagtt tgatcttgtc tggctctatg agagtgggtg gcacatgccg aacaaacaga 660
 agtttghtaag cgagctggca cgcgtcgag ctccaggagc aactatcatc atcgtgacct 720
 ggtgccatag gaacctcgcg ccgtcggagg actcactgaa acctgacgag ctgaatcttt 780
 tgaaaaagat ttgtgatgca tattacctcc cggattgttg ctgcgccctc gattatgtca 840
 agattgccga gtcattgtct cttgaggata tcaaaacggc cgactggtca gaaaacgtgg 900
 cccggttctg gcctgctgtc atccaatcag cactgacatg gaaaggcctc acttctctac 960
 taaggagtgg atggaagacg ataaagggag cactggtgat gcctctcatg atccaaggct 1020
 acaagaaagg cctcattaag ttcagcatca tcacctgccg caaaccccaa gcagccatag 1080
 aaggagaacc tgaggccgca tcgccagtg tagaatagaa cccatgtgat tggaatagac 1140
 tcggcttgcg gtcgcctcgt agctgaataa ttttgtgtta ccgtgcctct ctatctgcaa 1200
 ctggaagtgg cataggaaag tggttcctaa agcaaaaaaa aaaaaaaaaa aaaaaaa 1257

<210> 30
 <211> 366

<212> PRT
 <213> Triticum aestivum

<220>
 <221> UNSURE
 <222> (5)

<220>
 <221> UNSURE
 <222> (45)

<400> 30
 Met Ala Asn Ser Xaa Arg Pro Ala Pro Leu Thr Pro Leu His Arg Leu
 1 5 10 15
 Asp Ala Ala Pro Pro Pro Arg Pro Ser Leu Gly His Ala Ala Arg Pro
 20 25 30
 Val Pro Arg Pro Val Leu Pro Leu Leu Pro Ala Arg Xaa Leu Arg Ala
 35 40 45
 Pro Asp Gly Val Val Asp Asp Arg Gly Pro Gly Asp Ala Ala Pro Pro
 50 55 60
 Gly Leu Lys Glu Gly Ile Ala Gly Leu Tyr Asp Glu Ser Ser Gly Leu
 65 70 75 80
 Trp Glu Ser Ile Trp Gly Glu His Met His His Gly Phe Tyr Asp Ser
 85 90 95
 Gly Glu Ala Ala Ser Met Ser Asp His Arg Arg Ala Gln Ile Arg Met
 100 105 110
 Ile Glu Glu Ala Leu Ala Phe Ala Ala Val Pro Asp Asp Pro Thr Asn
 115 120 125
 Lys Pro Lys Thr Ile Val Asp Val Gly Cys Gly Ile Gly Gly Ser Ser
 130 135 140
 Arg Tyr Leu Ala Asn Lys Tyr Gly Ala Gln Cys Ser Gly Ile Thr Leu
 145 150 155 160
 Ser Pro Val Gln Ala Glu Arg Gly Asn Ala Leu Ala Ala Ala Gln Gly
 165 170 175
 Leu Ser Asp Lys Ala Ser Phe Gln Val Ala Asp Ala Leu Glu Gln Pro
 180 185 190
 Phe Pro Asp Gly Gln Phe Asp Leu Val Trp Ser Met Glu Ser Gly Glu
 195 200 205
 His Met Pro Asn Lys Gln Lys Phe Val Ser Glu Leu Ala Arg Val Ala
 210 215 220
 Ala Pro Gly Ala Thr Ile Ile Ile Val Thr Trp Cys His Arg Asn Leu
 225 230 235 240
 Ala Pro Ser Glu Asp Ser Leu Lys Pro Asp Glu Leu Asn Leu Leu Lys
 245 250 255

Lys Ile Cys Asp Ala Tyr Tyr Leu Pro Asp Trp Cys Ser Pro Ser Asp
260 265 270

Tyr Val Lys Ile Ala Glu Ser Leu Ser Leu Glu Asp Ile Lys Thr Ala
275 280 285

Asp Trp Ser Glu Asn Val Ala Pro Phe Trp Pro Ala Val Ile Gln Ser
290 295 300

Ala Leu Thr Trp Lys Gly Leu Thr Ser Leu Leu Arg Ser Gly Trp Lys
305 310 315 320

Thr Ile Lys Gly Ala Leu Val Met Pro Leu Met Ile Gln Gly Tyr Lys
325 330 335

Lys Gly Leu Ile Lys Phe Ser Ile Ile Thr Cys Arg Lys Pro Gln Ala
340 345 350

Ala Ile Glu Gly Glu Pro Glu Ala Ala Ser Pro Ser Val Glu
355 360 365

<210> 31
<211> 1605
<212> DNA
<213> Catalpa sp.

<400> 31
gcacgaggca ccacttcacc accaccacca ccaccaccac caccacaact gcttcaccg 60
ccgcagagtc actctatcag ttaagatcaa agcaacaagt tcagaatccc aaacaatggg 120
caagcagacg accacttcgc ccaccgccgc ggacgggtcc aaagatgcgc atgcagaatt 180
caagctgggtg ggcttcaaga atttcgtcag gaccaacccc aagtcgcacc acttctgcgt 240
ccaccgcttc caccatatag agttctggtg cggcgacgcc accaacaccg ccaagcgctt 300
ctcttggggc ctccggtatgc ccctcgtcgc caaatcggat ctttccactg gaaactccgc 360
tcatgcctcg tatcttcttc ggtctggcga actcaacttc ctcttcacga gcccttactc 420
tccttcaatc tcgcgccctt cctccgccgc catccccagt ttctccttct ccacctacca 480
gtctttttacc tcttcccatg gcctcgctgt tcgtgcgggtg gctattcagg tcgattcggc 540
cttttcggct tactctgctt ccatttcccg cggcgccaaa cccgtgtccg caccgattct 600
tttatctgac aacaagactg ccattgcgga ggttcattta tatggagact cagtgttgcg 660
attcgtgagc tatgggtgata atgggacagg cccagatgga tggttcttgc cgggctttga 720
gcctgtggat gatcagatgt cttataaaga atttgattat gggattagaa ggctagatca 780
tgctgtagga aatgtgcccg aactcgggtc agttgtggat tacttgaaaa aattcacagg 840
gtttcatgaa tttgcagagt ttacttcaga ggatgtggga acagcagaaa gtggattgaa 900
ttctatgggt ttagcgaaca acaatgaaaa tgtgttggtt cctctgaacg aaccgggtgt 960
tggaaccaag aggaagagcc agattcagac ttatttggag cacaatgaag ggccagggtg 1020
acagcatttg gcattagtga gtgaggatat ctttaacaca ttaagggaaa tgagaaagag 1080
gagtggagtt gggggattcg agttcatgcc ttgcctccg cttacttatt acaagaatct 1140
caagaacaga gctggagatg tgctgaggga tgagcagatt gaggagtgtg agaagttggg 1200
gatcttgggt gacagggatg atcaggggac tttgcttcag attttcacca agcctgtggg 1260
tgataggcca acgctattca tagagatcat tcagagaatc ggggtgcagc tcaaagacga 1320
acaaggaaag ctctaccaga agagtgggtt tggaggattt ggaaagggca acttctccga 1380
actcttcaaa tccatcgaag aatacgagaa aatgctcgaa gcaaaagcaag tcaactgaaac 1440
agegtcggcc tgagttctga gtccttctta ctgtgttgta gatatgttga tgaaccaatg 1500
tcctgtcggg acataggttg ttcttatgct gtactaaact gtagtgtgaca agaagtttta 1560
cttaataata tatcgtactt tctataaaaa aaaaaaaaaa aaaaa 1605

<210> 32
<211> 445
<212> PRT
<213> Catalpa sp.

<400> 32
 Met Gly Lys Gln Thr Thr Thr Ser Ala Thr Ala Ala Asp Gly Ser Lys
 1 5 10 15
 Asp Ala His Ala Glu Phe Lys Leu Val Gly Phe Lys Asn Phe Val Arg
 20 25 30
 Thr Asn Pro Lys Ser Asp His Phe Cys Val His Arg Phe His His Ile
 35 40 45
 Glu Phe Trp Cys Gly Asp Ala Thr Asn Thr Ala Lys Arg Phe Ser Trp
 50 55 60
 Gly Leu Gly Met Pro Leu Val Ala Lys Ser Asp Leu Ser Thr Gly Asn
 65 70 75 80
 Ser Ala His Ala Ser Tyr Leu Leu Arg Ser Gly Glu Leu Asn Phe Leu
 85 90 95
 Phe Thr Ser Pro Tyr Ser Pro Ser Ile Ser Ala Pro Ser Ser Ala Ala
 100 105 110
 Ile Pro Ser Phe Ser Phe Ser Thr Tyr Gln Ser Phe Thr Ser Ser His
 115 120 125
 Gly Leu Ala Val Arg Ala Val Ala Ile Gln Val Asp Ser Ala Phe Ser
 130 135 140
 Ala Tyr Ser Ala Ser Ile Ser Arg Gly Ala Lys Pro Val Ser Ala Pro
 145 150 155 160
 Ile Leu Leu Ser Asp Asn Lys Thr Ala Ile Ala Glu Val His Leu Tyr
 165 170 175
 Gly Asp Ser Val Leu Arg Phe Val Ser Tyr Gly Asp Asn Gly Thr Gly
 180 185 190
 Pro Asp Gly Trp Phe Leu Pro Gly Phe Glu Pro Val Asp Asp Gln Met
 195 200 205
 Ser Tyr Lys Glu Leu Asp Tyr Gly Ile Arg Arg Leu Asp His Ala Val
 210 215 220
 Gly Asn Val Pro Glu Leu Gly Pro Val Val Asp Tyr Leu Lys Lys Phe
 225 230 235 240
 Thr Gly Phe His Glu Phe Ala Glu Phe Thr Ser Glu Asp Val Gly Thr
 245 250 255
 Ala Glu Ser Gly Leu Asn Ser Met Val Leu Ala Asn Asn Asn Glu Asn
 260 265 270
 Val Leu Leu Pro Leu Asn Glu Pro Val Phe Gly Thr Lys Arg Lys Ser
 275 280 285
 Gln Ile Gln Thr Tyr Leu Glu His Asn Glu Gly Pro Gly Val Gln His
 290 295 300
 Leu Ala Leu Val Ser Glu Asp Ile Phe Asn Thr Leu Arg Glu Met Arg
 305 310 315 320

<400> 34
 Thr Arg Lys Ser Tyr Gly Leu Arg Arg Phe Asp His Val Val Gly Asn
 1 5 10 15

Val Pro Glu Leu Ala Pro Val Ala Ala Tyr Ile Ser Gly Phe Thr Gly
 20 25 30
 Phe His Glu Phe Ala Glu Phe Thr Ala Glu Asp Val Gly Thr Ala Glu
 35 40 45
 Ser Gly Leu Asn Ser Val Val Leu Ala Asn Asn Ala Glu Thr Val Leu
 50 55 60
 Leu Pro Leu Asn Glu Pro Val His Gly Thr Lys Arg Arg Ser Gln Ile
 65 70 75 80
 Gln Thr Tyr Leu Asp His His Gly Gly Pro Gly Val Gln His Ile Ala
 85 90 95
 Leu Ala Ser Asp Asp Val Leu Gly Thr Leu Arg Glu Met Arg Ala Arg
 100 105 110
 Ser Ala Met Gly Gly Phe Glu Phe Leu Ala Pro Pro Pro Pro Asn Tyr
 115 120 125
 Tyr Asp Gly Val Arg Arg Arg Ala Gly Asp Val Leu Ser Glu Glu Gln
 130 135 140
 Ile Asn Glu Cys Gln Glu Leu Gly Val Leu Val Asp Arg Asp Asp Gln
 145 150 155 160
 Gly Val Leu Leu Gln Ile Phe Thr Lys Pro Val Gly Asp Arg Pro Thr
 165 170 175
 Phe Phe Leu Glu Met Ile Gln Arg Ile Gly Cys Met Glu Lys Asp Glu
 180 185 190
 Ser Gly Gln Glu Tyr Gln Lys Gly Gly Cys Gly Gly Phe Gly Lys Gly
 195 200 205
 Asn Phe Ser Glu Leu Phe Lys Ser Ile Glu Glu Tyr Glu Lys Ser Leu
 210 215 220
 Glu Ala Lys Gln Ala Pro Thr Val Gln Gly Ser
 225 230 235

<210> 35
 <211> 1550
 <212> DNA
 <213> Glycine max

<400> 35
 tcacaccaca ccaatgccaa taccatgtg caacgaaatt caagcccaag cccaagccca 60
 agcccaagcc caacctgggt ttaagctcgt cggtttcaaa aacttcgtcc gaaccaatcc 120
 taagtccgac cgctttcaag tcaaccgctt ccaccacatc gagttctggt gcaccgatgc 180
 caccaacgcc tctcgccgat tctcttgggg acttggaatg cctattgttg caaaatctga 240
 tctctccacc ggaaaccaa tccacgcctc ctacctctc cgctccggcg acctctctt 300
 cctcttctcc gctccttact ctcectctct ctccgcggc tctcgcgtg cctcctccgc 360
 ctccattccc agtttcgacg ccgccacctg ccttgccctc gctgccaac acggcttcgg 420
 cgtcgcgcc atcgccttgg aagtcgccga cgcggaagcc gctttcagcg ccagcgtcgc 480
 gaaaggagcc gagccggcgt cgcgcgggt tctcgtcgac gatcgaccg gcttcgcgga 540
 ggtgcgcctc tacggcgacg tgggtgctcg ctacgtcagc tacaaggacg ccgcgcgcga 600
 ggccgcacac gcagatccgt cgcggtggtt cctgcgggga ttcgaggccg cggcgtcgtc 660
 gtcttcgttt ccggagctgg actacgggat ccggcggtg gaccacgccg tcgggaacgt 720


```

tccggagctg ggcggggcgg tgaggtagctt gaaaggcttc agcggattcc acgagttcgc 780
ggagttcacc gcggaggacg tgggaacgag cgagagcggg ttgaactcgg tggttctggc 840
gaacaactcg gagacggtgt tgctgccgct gaacgagccg gtttacggaa cgaagaggaa 900
gagccagatt gagacgtatt tggaacacaa cgaagggtgt ggtgtgcagc accttgcgct 960
tgttactcac gacatcttca ccacactgag agagatgaga aagcgaagtt tccttgggtg 1020
atattgagttc atgccttctc ctccctccac ctattacgcc aacctccaca accgtgccgc 1080
tgatgtgttg accgttgacc agattaagca gtgtgaggag cttgggattc ttgttgacag 1140
agatgatcag ggcactctgc ttcagatttt caccaagcct gttggggaca ggccaacgat 1200
attcatagag ataattcaga ggatcgggtg catggtggag gatgaggaag ggaaggtgta 1260
ccagaagggt gcatgtgggg gttttgggaa aggcaatttt tctgagcttt tcaaattccat 1320
tgaagaatat gagaagactt tggaagctaa aagaaccgag taagcacatt ggaagaacac 1380
aaataactct ttgttgaaat gattaatgag gaatcaatgt ggcatagggg gttttatactc 1440
tataatacat agaattacaa tgatagtgtc ctcccttgta tgaaaatgaa atcacagaaa 1500
cttttatgga tagtatTTTT ctattaaaaa aaaaaaaaaa aaaaaaaaaa 1550

```

<210> 36
 <211> 449
 <212> PRT
 <213> Glycine max

```

<400> 36
Met Pro Ile Pro Met Cys Asn Glu Ile Gln Ala Gln Ala Gln Ala Gln
  1              5              10              15

Ala Gln Ala Gln Pro Gly Phe Lys Leu Val Gly Phe Lys Asn Phe Val
      20              25              30

Arg Thr Asn Pro Lys Ser Asp Arg Phe Gln Val Asn Arg Phe His His
      35              40              45

Ile Glu Phe Trp Cys Thr Asp Ala Thr Asn Ala Ser Arg Arg Phe Ser
      50              55              60

Trp Gly Leu Gly Met Pro Ile Val Ala Lys Ser Asp Leu Ser Thr Gly
      65              70              75              80

Asn Gln Ile His Ala Ser Tyr Leu Leu Arg Ser Gly Asp Leu Ser Phe
      85              90              95

Leu Phe Ser Ala Pro Tyr Ser Pro Ser Leu Ser Ala Gly Ser Ser Ala
      100             105             110

Ala Ser Ser Ala Ser Ile Pro Ser Phe Asp Ala Ala Thr Cys Leu Ala
      115             120             125

Phe Ala Ala Lys His Gly Phe Gly Val Arg Ala Ile Ala Leu Glu Val
      130             135             140

Ala Asp Ala Glu Ala Ala Phe Ser Ala Ser Val Ala Lys Gly Ala Glu
      145             150             155             160

Pro Ala Ser Pro Pro Val Leu Val Asp Asp Arg Thr Gly Phe Ala Glu
      165             170             175

Val Arg Leu Tyr Gly Asp Val Val Leu Arg Tyr Val Ser Tyr Lys Asp
      180             185             190

Ala Ala Pro Gln Ala Pro His Ala Asp Pro Ser Arg Trp Phe Leu Pro
      195             200             205

```

Gly Phe Glu Ala Ala Ala Ser Ser Ser Ser Phe Pro Glu Leu Asp Tyr
 210 215 220
 Gly Ile Arg Arg Leu Asp His Ala Val Gly Asn Val Pro Glu Leu Ala
 225 230 235 240
 Pro Ala Val Arg Tyr Leu Lys Gly Phe Ser Gly Phe His Glu Phe Ala
 245 250 255
 Glu Phe Thr Ala Glu Asp Val Gly Thr Ser Glu Ser Gly Leu Asn Ser
 260 265 270
 Val Val Leu Ala Asn Asn Ser Glu Thr Val Leu Leu Pro Leu Asn Glu
 275 280 285
 Pro Val Tyr Gly Thr Lys Arg Lys Ser Gln Ile Glu Thr Tyr Leu Glu
 290 295 300
 His Asn Glu Gly Ala Gly Val Gln His Leu Ala Leu Val Thr His Asp
 305 310 315 320
 Ile Phe Thr Thr Leu Arg Glu Met Arg Lys Arg Ser Phe Leu Gly Gly
 325 330 335
 Phe Glu Phe Met Pro Ser Pro Pro Pro Thr Tyr Tyr Ala Asn Leu His
 340 345 350
 Asn Arg Ala Ala Asp Val Leu Thr Val Asp Gln Ile Lys Gln Cys Glu
 355 360 365
 Glu Leu Gly Ile Leu Val Asp Arg Asp Asp Gln Gly Thr Leu Leu Gln
 370 375 380
 Ile Phe Thr Lys Pro Val Gly Asp Arg Pro Thr Ile Phe Ile Glu Ile
 385 390 395 400
 Ile Gln Arg Ile Gly Cys Met Val Glu Asp Glu Glu Gly Lys Val Tyr
 405 410 415
 Gln Lys Gly Ala Cys Gly Gly Phe Gly Lys Gly Asn Phe Ser Glu Leu
 420 425 430
 Phe Lys Ser Ile Glu Glu Tyr Glu Lys Thr Leu Glu Ala Lys Arg Thr
 435 440 445

Ala

<210> 37
 <211> 1614
 <212> DNA
 <213> Triticum aestivum

<400> 37
 gcacgagcaa gaagcgaaca cacaccatgc cgcccacccc caccaccccc gcagccaccg 60
 gcgccgccgc ggtgacgccg gagcacgcgc gccgcgccg aatgggtccgc ttcaaccgcg 120
 gcagcgaccg ctccacacg ctgccttcc accacgtcga gttctggtgc gcggacgccg 180
 cctccgccgc cgcccgcttc gccttcgcgc tcggcgccgc gtcgccgcc aggtccgacc 240
 tctccacggg gaactccgtg cagcctccc agctgctccg ctggggcaac ctgccttcc 300
 tcttcacggc cccctacgcc aacggctgcg acgccgccac cgctccctg ccctccttct 360

```

ccgccgacgc cgcgcgccag ttctccgcgg accacggcct cgcggtgcgc tccatagcgc 420
tgccgcgtcgc ggacgctgcc gaggccttcc gcgccagcgt cgacgggggc gcgcgcccgc 480
ccttcagccc tgtggacctc ggccgcgggt tcggcttcgc ggaggtcgag ctctacggcg 540
acgtcgtgct ccgcttcgtc agccacccgg acggcaggga cgtgcccttc ttgccggggg 600
tcgagggcgt gagcaacca gacgccgtgg actacggcct gacgcggttc gaccacgtcg 660
tcggcaacgt cccggagctt gccccgcgcg cggcctacgt cgcggggttc acgggggttc 720
acgagttcgc cgagttcacg acggaggacg tgggcacggc cgagagcggg ctcaactcga 780
tggtgctcgc caacaactcg gagggcgtgc tgctgccgct caacgagccg gtgcacggca 840
ccaagcgccg gagccagata cagacgttcc tggaaacacca cggcggctcg ggcgtgcagc 900
acatcgccgt ggccagcagc gacgtgctca ggacgtcag ggagatgcgt gcgcgtcccg 960
ccatgggcgg cttcgacttc ctgccacccc cgctgccgaa gtactacgaa ggcgtgcggc 1020
gcatcgccgg ggatgtgctc tcggaggcgc agatcaagga atgccaggag ctgggggtgc 1080
tcgtcgacag ggacgaccaa ggggtgttgc taaaaatctt caccaagcca gtaggggaca 1140
ggccgacgtt gttcctggag atgatccaga ggatcgggtg catggagaag gacgagagag 1200
gggaagagta ccagaagggt ggctgcggcg ggttcggcaa aggcaacttc tccgagctgt 1260
tcaagtccat tgaagattac gagaagtccc ttgaagccaa gcaatctgct gcagttcagg 1320
gatcatagga tagaagctgg agctggagga gctgatccag tactttgtat caggtctcat 1380
ggagcaaaag aaaatgatgt tgtttgtaag atcgcgcgcg caattatgtc cgatgttata 1440
attggtgaag ctgaagacag atgtatccta tgtatgatgg gtgtaataga tggtagaggg 1500
ggctcggctc acacatgaac aaaatgtact gttggcattg ttgtataatc ttgcttgcaa 1560
gtaaaataaa gaagaaccga ttttgagttc tgcataaaaa aaaaaaaaaa aaaa .1614

```

<210> 38
 <211> 433
 <212> PRT
 <213> Triticum aestivum

<400> 38
 Met Pro Pro Thr Pro Thr Thr Pro Ala Ala Thr Gly Ala Ala Ala Val
 1 5 10 15
 Thr Pro Glu His Ala Arg Pro Arg Arg Met Val Arg Phe Asn Pro Arg
 20 25 30
 Ser Asp Arg Phe His Thr Leu Ala Phe His His Val Glu Phe Trp Cys
 35 40 45
 Ala Asp Ala Ala Ser Ala Ala Gly Arg Phe Ala Phe Ala Leu Gly Ala
 50 55 60
 Pro Leu Ala Ala Arg Ser Asp Leu Ser Thr Gly Asn Ser Val His Ala
 65 70 75 80
 Ser Gln Leu Leu Arg Ser Gly Asn Leu Ala Phe Leu Phe Thr Ala Pro
 85 90 95
 Tyr Ala Asn Gly Cys Asp Ala Ala Thr Ala Ser Leu Pro Ser Phe Ser
 100 105 110
 Ala Asp Ala Ala Arg Gln Phe Ser Ala Asp His Gly Leu Ala Val Arg
 115 120 125
 Ser Ile Ala Leu Arg Val Ala Asp Ala Ala Glu Ala Phe Arg Ala Ser
 130 135 140
 Val Asp Gly Gly Ala Arg Pro Ala Phe Ser Pro Val Asp Leu Gly Arg
 145 150 155 160
 Gly Phe Gly Phe Ala Glu Val Glu Leu Tyr Gly Asp Val Val Leu Arg
 165 170 175

Phe Val Ser His Pro Asp Gly Arg Asp Val Pro Phe Leu Pro Gly Phe
 180 185 190
 Glu Gly Val Ser Asn Pro Asp Ala Val Asp Tyr Gly Leu Thr Arg Phe
 195 200 205
 Asp His Val Val Gly Asn Val Pro Glu Leu Ala Pro Ala Ala Ala Tyr
 210 215 220
 Val Ala Gly Phe Thr Gly Phe His Glu Phe Ala Glu Phe Thr Thr Glu
 225 230 235 240
 Asp Val Gly Thr Ala Glu Ser Gly Leu Asn Ser Met Val Leu Ala Asn
 245 250 255
 Asn Ser Glu Gly Val Leu Leu Pro Leu Asn Glu Pro Val His Gly Thr
 260 265 270
 Lys Arg Arg Ser Gln Ile Gln Thr Phe Leu Glu His His Gly Gly Ser
 275 280 285
 Gly Val Gln His Ile Ala Val Ala Ser Ser Asp Val Leu Arg Thr Leu
 290 295 300
 Arg Glu Met Arg Ala Arg Ser Ala Met Gly Gly Phe Asp Phe Leu Pro
 305 310 315 320
 Pro Pro Leu Pro Lys Tyr Tyr Glu Gly Val Arg Arg Ile Ala Gly Asp
 325 330 335
 Val Leu Ser Glu Ala Gln Ile Lys Glu Cys Gln Glu Leu Gly Val Leu
 340 345 350
 Val Asp Arg Asp Asp Gln Gly Val Leu Leu Gln Ile Phe Thr Lys Pro
 355 360 365
 Val Gly Asp Arg Pro Thr Leu Phe Leu Glu Met Ile Gln Arg Ile Gly
 370 375 380
 Cys Met Glu Lys Asp Glu Arg Gly Glu Glu Tyr Gln Lys Gly Gly Cys
 385 390 395 400
 Gly Gly Phe Gly Lys Gly Asn Phe Ser Glu Leu Phe Lys Ser Ile Glu
 405 410 415
 Asp Tyr Glu Lys Ser Leu Glu Ala Lys Gln Ser Ala Ala Val Gln Gly
 420 425 430

Ser

<210> 39
 <211> 317
 <212> PRT
 <213> Synechocystis sp.

<400> 39
 Met Val Tyr His Val Arg Pro Lys His Ala Leu Phe Leu Ala Phe Tyr
 1 5 10 15

Cys Tyr Phe Ser Leu Leu Thr Met Ala Ser Ala Thr Ile Ala Ser Ala
 20 25 30
 Asp Leu Tyr Glu Lys Ile Lys Asn Phe Tyr Asp Asp Ser Ser Gly Leu
 35 40 45
 Trp Glu Asp Val Trp Gly Glu His Met His His Gly Tyr Tyr Gly Pro
 50 55 60
 His Gly Thr Tyr Arg Ile Asp Arg Arg Gln Ala Gln Ile Asp Leu Ile
 65 70 75 80
 Lys Glu Leu Leu Ala Trp Ala Val Pro Gln Asn Ser Ala Lys Pro Arg
 85 90 95
 Lys Ile Leu Asp Leu Gly Cys Gly Ile Gly Gly Ser Ser Leu Tyr Leu
 100 105 110
 Ala Gln Gln His Gln Ala Glu Val Met Gly Ala Ser Leu Ser Pro Val
 115 120 125
 Gln Val Glu Arg Ala Gly Glu Arg Ala Arg Ala Leu Gly Leu Gly Ser
 130 135 140
 Thr Cys Gln Phe Gln Val Ala Asn Ala Leu Asp Leu Pro Phe Ala Ser
 145 150 155 160
 Asp Ser Phe Asp Trp Val Trp Ser Leu Glu Ser Gly Glu His Met Pro
 165 170 175
 Asn Lys Ala Gln Phe Leu Gln Glu Ala Trp Arg Val Leu Lys Pro Gly
 180 185 190
 Gly Arg Leu Ile Leu Ala Thr Trp Cys His Arg Pro Ile Asp Pro Gly
 195 200 205
 Asn Gly Pro Leu Thr Ala Asp Glu Arg Arg His Leu Gln Ala Ile Tyr
 210 215 220
 Asp Val Tyr Cys Leu Pro Tyr Val Val Ser Leu Pro Asp Tyr Glu Ala
 225 230 235 240
 Ile Ala Arg Glu Cys Gly Phe Gly Glu Ile Lys Thr Ala Asp Trp Ser
 245 250 255
 Val Ala Val Ala Pro Phe Trp Asp Arg Val Ile Glu Ser Ala Phe Asp
 260 265 270
 Pro Arg Val Leu Trp Ala Leu Gly Gln Ala Gly Pro Lys Ile Ile Asn
 275 280 285
 Ala Ala Leu Cys Leu Arg Leu Met Lys Trp Gly Tyr Glu Arg Gly Leu
 290 295 300
 Val Arg Phe Gly Leu Leu Thr Gly Ile Lys Pro Leu Val
 305 310 315
 <210> 40
 <211> 348

<212> PRT

<213> Arabidopsis thaliana

<400> 40

Met Lys Ala Thr Leu Ala Ala Pro Ser Ser Leu Thr Ser Leu Pro Tyr
 1 5 10 15
 Arg Thr Asn Ser Ser Phe Gly Ser Lys Ser Ser Leu Leu Phe Arg Ser
 20 25 30
 Pro Ser Ser Ser Ser Ser Val Ser Met Thr Thr Thr Arg Gly Asn Val
 35 40 45
 Ala Val Ala Ala Ala Ala Thr Ser Thr Glu Ala Leu Arg Lys Gly Ile
 50 55 60
 Ala Glu Phe Tyr Asn Glu Thr Ser Gly Leu Trp Glu Glu Ile Trp Gly
 65 70 75 80
 Asp His Met His His Gly Phe Tyr Asp Pro Asp Ser Ser Val Gln Leu
 85 90 95
 Ser Asp Ser Gly His Lys Glu Ala Gln Ile Arg Met Ile Glu Glu Ser
 100 105 110
 Leu Arg Phe Ala Gly Val Thr Asp Glu Glu Glu Glu Lys Lys Ile Lys
 115 120 125
 Lys Val Val Asp Val Gly Cys Gly Ile Gly Gly Ser Ser Arg Tyr Leu
 130 135 140
 Ala Ser Lys Phe Gly Ala Glu Cys Ile Gly Ile Thr Leu Ser Pro Val
 145 150 155 160
 Gln Ala Lys Arg Ala Asn Asp Leu Ala Ala Ala Gln Ser Leu Ser His
 165 170 175
 Lys Ala Ser Phe Gln Val Ala Asp Ala Leu Asp Gln Pro Phe Glu Asp
 180 185 190
 Gly Lys Phe Asp Leu Val Trp Ser Met Glu Ser Gly Glu His Met Pro
 195 200 205
 Asp Lys Ala Lys Phe Val Lys Glu Leu Val Arg Val Ala Ala Pro Gly
 210 215 220
 Gly Arg Ile Ile Ile Val Thr Trp Cys His Arg Asn Leu Ser Ala Gly
 225 230 235 240
 Glu Glu Ala Leu Gln Pro Trp Glu Gln Asn Ile Leu Asp Lys Ile Cys
 245 250 255
 Lys Thr Phe Tyr Leu Pro Ala Trp Cys Ser Thr Asp Asp Tyr Val Asn
 260 265 270
 Leu Leu Gln Ser His Ser Leu Gln Asp Ile Lys Cys Ala Asp Trp Ser
 275 280 285
 Glu Asn Val Ala Pro Phe Trp Pro Ala Val Ile Arg Thr Ala Leu Thr
 290 295 300

Trp Lys Gly Leu Val Ser Leu Leu Arg Ser Gly Met Lys Ser Ile Lys
305 310 315 320

Gly Ala Leu Thr Met Pro Leu Met Ile Glu Gly Tyr Lys Lys Gly Val
325 330 335

Ile Lys Phe Gly Ile Ile Thr Cys Gln Lys Pro Leu
340 345

<210> 41

<211> 434

<212> PRT

<213> Hordeum vulgare

<400> 41

Met Pro Pro Thr Pro Thr Thr Pro Ala Ala Thr Gly Ala Ala Ala Ala
1 5 10 15

Val Thr Pro Glu His Ala Arg Pro His Arg Met Val Arg Phe Asn Pro
20 25 30

Arg Ser Asp Arg Phe His Thr Leu Ser Phe His His Val Glu Phe Trp
35 40 45

Cys Ala Asp Ala Ala Ser Ala Ala Gly Arg Phe Ala Phe Ala Leu Gly
50 55 60

Ala Pro Leu Ala Ala Arg Ser Asp Leu Ser Thr Gly Asn Ser Ala His
65 70 75 80

Ala Ser Gln Leu Leu Arg Ser Gly Ser Leu Ala Phe Leu Phe Thr Ala
85 90 95

Pro Tyr Ala Asn Gly Cys Asp Ala Ala Thr Ala Ser Leu Pro Ser Phe
100 105 110

Ser Ala Asp Ala Ala Arg Arg Phe Ser Ala Asp His Gly Ile Ala Val
115 120 125

Arg Ser Val Ala Leu Arg Val Ala Asp Ala Ala Glu Ala Phe Arg Ala
130 135 140

Ser Arg Arg Arg Gly Ala Arg Pro Ala Phe Ala Pro Val Asp Leu Gly
145 150 155 160

Arg Gly Phe Ala Phe Ala Glu Val Glu Leu Tyr Gly Asp Val Val Leu
165 170 175

Arg Phe Val Ser His Pro Asp Gly Thr Asp Val Pro Phe Leu Pro Gly
180 185 190

Phe Glu Gly Val Thr Asn Pro Asp Ala Val Asp Tyr Gly Leu Thr Arg
195 200 205

Phe Asp His Val Val Gly Asn Val Pro Glu Leu Ala Pro Ala Ala Ala
210 215 220

Tyr Ile Ala Gly Phe Thr Gly Phe His Glu Phe Ala Glu Phe Thr Ala
225 230 235 240

Glu Asp Val Gly Thr Thr Glu Ser Gly Leu Asn Ser Val Val Leu Ala
 245 250 255
 Asn Asn Ser Glu Gly Val Leu Leu Pro Leu Asn Glu Pro Val His Gly
 260 265 270
 Thr Lys Arg Arg Ser Gln Ile Gln Thr Phe Leu Glu His His Gly Gly
 275 280 285
 Pro Gly Val Gln His Ile Ala Val Ala Ser Ser Asp Val Leu Arg Thr
 290 295 300
 Leu Arg Lys Met Arg Ala Arg Ser Ala Met Gly Gly Phe Asp Phe Leu
 305 310 315 320
 Pro Pro Pro Leu Pro Lys Tyr Tyr Glu Gly Val Arg Arg Leu Ala Gly
 325 330 335
 Asp Val Leu Ser Glu Ala Gln Ile Lys Glu Cys Gln Glu Leu Gly Val
 340 345 350
 Leu Val Asp Arg Asp Asp Gln Gly Val Leu Leu Gln Ile Phe Thr Lys
 355 360 365
 Pro Val Gly Asp Arg Pro Thr Leu Phe Leu Glu Met Ile Gln Arg Ile
 370 375 380
 Gly Cys Met Glu Lys Asp Glu Arg Gly Glu Glu Tyr Gln Lys Gly Gly
 385 390 395 400
 Cys Gly Gly Phe Gly Lys Gly Asn Phe Ser Glu Leu Phe Lys Ser Ile
 405 410 415
 Glu Asp Tyr Glu Lys Ser Leu Glu Ala Lys Gln Ser Ala Ala Val Gln
 420 425 430

Gly Ser

<210> 42
 <211> 442
 <212> PRT
 <213> Daucus carota

<400> 42
 Met Gly Lys Lys Gln Ser Glu Ala Glu Ile Leu Ser Ser Asn Ser Ser
 1 5 10 15
 Asn Thr Ser Pro Ala Thr Phe Lys Leu Val Gly Phe Asn Asn Phe Val
 20 25 30
 Arg Ala Asn Pro Lys Ser Asp His Phe Ala Val Lys Arg Phe His His
 35 40 45
 Ile Glu Phe Trp Cys Gly Asp Ala Thr Asn Thr Ser Arg Arg Phe Ser
 50 55 60
 Trp Gly Leu Gly Met Pro Leu Val Ala Lys Ser Asp Leu Ser Thr Gly
 65 70 75 80

Asn Ser Val His Ala Ser Tyr Leu Val Arg Ser Ala Asn Leu Ser Phe
 85 90 95
 Val Phe Thr Ala Pro Tyr Ser Pro Ser Thr Thr Thr Ser Ser Gly Ser
 100 105 110
 Ala Ala Ile Pro Ser Phe Ser Ala Ser Gly Phe His Ser Phe Ala Ala
 115 120 125
 Lys His Gly Leu Ala Val Arg Ala Ile Ala Leu Glu Val Ala Asp Val
 130 135 140
 Ala Ala Ala Phe Glu Ala Ser Val Ala Arg Gly Ala Arg Pro Ala Ser
 145 150 155 160
 Ala Pro Val Glu Leu Asp Asp Gln Ala Trp Leu Ala Glu Val Glu Leu
 165 170 175
 Tyr Gly Asp Val Val Leu Arg Phe Val Ser Phe Gly Arg Glu Glu Gly
 180 185 190
 Leu Phe Leu Pro Gly Phe Glu Ala Val Glu Gly Thr Ala Ser Phe Pro
 195 200 205
 Asp Leu Asp Tyr Gly Ile Arg Arg Leu Asp His Ala Val Gly Asn Val
 210 215 220
 Thr Glu Leu Gly Pro Val Val Glu Tyr Ile Lys Gly Phe Thr Gly Phe
 225 230 235 240
 His Glu Phe Ala Glu Phe Thr Ala Glu Asp Val Gly Thr Leu Glu Ser
 245 250 255
 Gly Leu Asn Ser Val Val Leu Ala Asn Asn Glu Glu Met Val Leu Leu
 260 265 270
 Pro Leu Asn Glu Pro Val Tyr Gly Thr Lys Arg Lys Ser Gln Ile Gln
 275 280 285
 Thr Tyr Leu Glu His Asn Glu Gly Ala Gly Val Gln His Leu Ala Leu
 290 295 300
 Val Ser Glu Asp Ile Phe Arg Thr Leu Arg Glu Met Arg Lys Arg Ser
 305 310 315 320
 Cys Leu Gly Gly Phe Glu Phe Met Pro Ser Pro Pro Pro Thr Tyr Tyr
 325 330 335
 Lys Asn Leu Lys Asn Arg Val Gly Asp Val Leu Ser Asp Glu Gln Ile
 340 345 350
 Lys Glu Cys Glu Asp Leu Gly Ile Leu Val Asp Arg Asp Asp Gln Gly
 355 360 365
 Thr Leu Leu Gln Ile Phe Thr Lys Pro Val Gly Asp Arg Pro Thr Leu
 370 375 380
 Phe Ile Glu Ile Ile Gln Arg Val Gly Cys Met Leu Lys Asp Asp Ala
 385 390 395 400

Gly Gln Met Tyr Gln Lys Gly Gly Cys Gly Gly Phe Gly Lys Gly Asn
 405 410 415

Phe Ser Glu Leu Phe Lys Ser Ile Glu Glu Tyr Glu Lys Thr Leu Glu
 420 425 430

Ala Lys Gln Ile Thr Gly Ser Ala Ala Ala
 435 440

<210> 43
 <211> 445
 <212> PRT
 <213> Arabidopsis thaliana

<400> 43
 Met Gly His Gln Asn Ala Ala Val Ser Glu Asn Gln Asn His Asp Asp
 1 5 10 15

Gly Ala Ala Ser Ser Pro Gly Phe Lys Leu Val Gly Phe Ser Lys Phe
 20 25 30

Val Arg Lys Asn Pro Lys Ser Asp Lys Phe Lys Val Lys Arg Phe His
 35 40 45

His Ile Glu Phe Trp Cys Gly Asp Ala Thr Asn Val Ala Arg Arg Phe
 50 55 60

Ser Trp Gly Leu Gly Met Arg Phe Ser Ala Lys Ser Asp Leu Ser Thr
 65 70 75 80

Gly Asn Met Val His Ala Ser Tyr Leu Leu Thr Ser Gly Asp Leu Arg
 85 90 95

Phe Leu Phe Thr Ala Pro Tyr Ser Pro Ser Leu Ser Ala Gly Glu Ile
 100 105 110

Lys Pro Thr Thr Thr Ala Ser Ile Pro Ser Phe Asp His Gly Ser Cys
 115 120 125

Arg Ser Phe Phe Ser Ser His Gly Leu Gly Val Arg Ala Val Ala Ile
 130 135 140

Glu Val Glu Asp Ala Glu Ser Ala Phe Ser Ile Ser Val Ala Asn Gly
 145 150 155 160

Ala Ile Pro Ser Ser Pro Pro Ile Val Leu Asn Glu Ala Val Thr Ile
 165 170 175

Ala Glu Val Lys Leu Tyr Gly Asp Val Val Leu Arg Tyr Val Ser Tyr
 180 185 190

Lys Ala Glu Asp Thr Glu Lys Ser Glu Phe Leu Pro Gly Phe Glu Arg
 195 200 205

Val Glu Asp Ala Ser Ser Phe Pro Leu Asp Tyr Gly Ile Arg Arg Leu
 210 215 220

Asp His Ala Val Gly Asn Val Pro Glu Leu Gly Pro Ala Leu Thr Tyr
 225 230 235 240

Val Ala Gly Phe Thr Gly Phe His Gln Phe Ala Glu Phe Thr Ala Asp
 245 250 255
 Asp Val Gly Thr Ala Glu Ser Gly Leu Asn Ser Ala Val Leu Ala Ser
 260 265 270
 Asn Asp Glu Met Val Leu Leu Pro Ile Asn Glu Pro Val His Gly Thr
 275 280 285
 Lys Arg Lys Ser Gln Ile Gln Thr Tyr Leu Glu His Asn Glu Gly Ala
 290 295 300
 Gly Leu Gln His Leu Ala Leu Met Ser Glu Asp Ile Phe Arg Thr Leu
 305 310 315 320
 Arg Glu Met Arg Lys Arg Ser Ser Ile Gly Gly Phe Asp Phe Met Pro
 325 330 335
 Ser Pro Pro Pro Thr Tyr Tyr Gln Asn Leu Lys Lys Arg Val Gly Asp
 340 345 350
 Val Leu Ser Asp Asp Gln Ile Lys Glu Cys Glu Glu Leu Gly Ile Leu
 355 360 365
 Val Asp Arg Asp Asp Gln Gly Thr Leu Leu Gln Ile Phe Thr Lys Pro
 370 375 380
 Leu Gly Asp Arg Pro Thr Ile Phe Ile Glu Ile Ile Gln Arg Val Gly
 385 390 395 400
 Cys Met Met Lys Asp Glu Glu Gly Lys Ala Tyr Gln Ser Gly Gly Cys
 405 410 415
 Gly Gly Phe Gly Lys Gly Asn Phe Ser Glu Leu Phe Lys Ser Ile Glu
 420 425 430
 Glu Tyr Glu Lys Thr Leu Glu Ala Lys Gln Leu Val Gly
 435 440 445